5 Implementation and Evaluation Plan

Establishing and implementing a successful Natural Resources Management Program (NRM Program) requires careful planning, sufficient funding, adequate staffing, and ongoing review and evaluation of program successes and failures. This section outlines the nuts and bolts of developing the NRM Program for the UH Management Areas on Mauna Kea and for implementing the management actions developed in Section 4. It also provides a methodology for evaluating the success of the program and for determining any need for changes in management strategies.

5.1 Implementation Plan

5.1.1 Programmatic Management Actions

Most of the goals, objectives, and actions of the NRM Program are presented in the component plans in Section 4. However, there are several management actions at the programmatic level that apply to either the overall NRM Program or that should be completed prior to enacting the actions in the various component plans. These programmatic management actions include:

- 1. Complete formal adoption of the NRMP by the University of Hawai'i. Initial implementation of the plan components can be undertaken by existing OMKM staff while the funding and staffing activities are in progress.
- 2. Establish a Natural Resources Management Program within the Office of Mauna Kea Management (OMKM).
 - a. Obtain sufficient funding to support the program (see Section 5.1.4).
 - b. Hire staff members. Although many of the management activities can be conducted with the aid of volunteers or in collaboration with other land management agencies, the minimum staffing needs for this program are one full-time Natural Resources Coordinator (NRC) and one or two full time field biologists. Support and administrative staff, such as secretaries and librarians, are already available at OMKM (see Section 5.1.2.1).
 - c. Contract out high-priority, field-intensive projects such as baseline inventory studies, to ensure they are conducted quickly (see Section 4.1).
- 3. Continue and increase the role of the Mauna Kea Management Board (MKMB) Environment Committee in the NRM Program.

The establishment of the MKMB and its associated committees has enabled the broader community to become directly involved in advising on the management of Mauna Kea. The committees have been useful in involving a range of interests to ensure a collaborative and informed management planning process. Each of the committees has its own mission, but there are few guidelines for how decisions are made, especially if there is no consensus or if the requisite expertise is not present. For example, the MKMB Environment Committee might more efficiently make defensible decisions if they had a protocol for evaluating the potential environmental impacts of proposed activities. The following activities are recommended to increase the decision-making power and participation in the NRM Program by the MKMB Environment Committee.

a. Review and approve (or request revisions to) management decisions, plans, and reports made by the NRC (via semi-annual meetings and as needed).

- i. Participate in annual and five-year reviews and revisions of Natural Resources Management Plan (NRMP). See Section 5.2.
- b. Review and approve or reject (with input from NRC and expert advice, as needed) research proposals from outside persons or agencies.
- c. Interact with other boards and committees (with participation of the NRC) to achieve agreement on management goals, objectives, and activities.¹
- d. Incorporate decisions from cross-board meetings into the NRMP (during revisions), and into daily activities conducted by the NRC. The NRC should keep detailed notes of decisions made during these board meetings for use during revision process, as it may be a year or more until the next NRMP revision cycle begins.
- 4. Continue to include the other advisory committees as integral parts of the NRM Program. The following committees and councils should meet annually (and as needed) with the NRC and the MKMB Environment Committee, to discuss natural resource management issues and strategies, including potential compatibility and conflicts with cultural issues:
 - a. Wēkiu Bug Committee
 - b. Kahu Kū Mauna
- 5. Continue to develop working relationships with federal and state agencies.

Currently there is no mechanism for integrated or coordinated management of Mauna Kea's natural resources (including lands outside of the UH Management Areas). No regular meetings are held between the governmental agencies with management responsibilities for Mauna Kea—in particular involving OMKM and the various divisions of DLNR. Significantly, because there is so little interaction between the various state and federal agencies responsible for the management of Mauna Kea, applicable rules and regulations in the UH Management Areas are little enforced. Development of coordinated management between state and federal agencies and OMKM is discussed in Section 5.1.3.

6. To increase community involvement in decision-making processes, conduct public outreach and education.

Encourage and seek out community participation, including Native Hawaiians, in educational and outreach efforts. Education and outreach activities are discussed further in Section 4.4. Education and outreach relating to natural resources will be conducted or overseen by the NRC, with input from the OMKM director and MKMB Environment Committee.²

7. Establish legal authority for OMKM to promulgate rules and enforce regulations on Mauna Kea properties leased by the University of Hawai'i.³

¹ As the advisory board for OMKM, the MKMB has responsibility for making final recommendations, based on input from committees including Kahu Kū Mauna and the MKMB Environment Committee. Potential conflicts between interests must be considered as this is done.

² The NRC will work closely with an education and outreach coordinator when this position is filled.

³ Although it is desirable that this action be completed prior to establishing the NRM Program, it is not a requirement. There are many management activities and projects that do not require regulatory authority, including baseline inventory, monitoring, research, education, and information management. Aspects of natural resources management that require regulatory authority are managing access, providing rangers with the authority to enforce laws, and establishing fines for rule-breaking. It is recommended that obtaining rule-making authority be completed concurrently with the establishment of the NRM Program, and that the program operate regardless of OMKM's ability to enforce regulations (see Section 1.4.2.3).

- 8. Manage access to the summit to reduce impacts to natural resources. Unrestricted access to the summit region can negatively impact resources. See Sections 3 and 4.2 for more information on the potential impacts of human use. Strategies for access management are presented in Sections 4.2 and 4.4.
- 9. Provide that the MKMB Environment Committee will have an active role in advising MKMB and OMKM with regard to natural resources when considering proposed future land uses (see Section 3.1.1.4) and ensuring the compatibility with recommendations in approved management plans, with the goal of protecting natural resources.

Although the UH Board of Regents and the President retain project approval and design review authority over all major developments within the UH Management Areas on Mauna Kea, OMKM, MKMB, and Kahu Kū Mauna are also charged with reviewing projects to ensure that they conform to the standards and guidelines set forth in the 2000 Master Plan. The 2000 Master Plan established a set of guidelines for project review and design, to ensure that proposed projects conform to and implement the concepts, themes, development standards and guidelines set forth in the plan (see Section XI, (Group 70 International 2000)). A Design Review Committee was established to interpret the guidelines and to ensure that proposed projects conform to the goals and objectives of the 2000 Master Plan and are consistent with the design guidelines.

The review process requires OMKM to work with other entities, including IfA, the University system, and DLNR. In addition to complying with federal, state and county rules and regulations, project proposers must be informed of additional conditions that OMKM may require. There is also a need, during the project design review process, for OMKM to provide clear facility planning guidelines to project proposers that address siting and design considerations, and to enforce them, so that proposed facilities result in minimal impacts to cultural and natural resources and the astronomical qualities of the MKSR. Many of these considerations have been developed in the 2000 Master Plan, and the MKMB Environment Committee should provide feedback and guidance on issues related to natural resources.

As specified in the 2000 Master Plan, each redevelopment or proposed new facility, including non-astronomy facilities, will undergo individual project reviews, that will include an environmental analysis pursuant to Chapter 343, Hawai'i Revised Statutes and a comprehensive analysis of the potential cultural impact. In general, the review process is applicable to any project involving any construction, installation or alteration upon any site, roadway, utility line, building, or other type of structure; any excavation, filling or change to surface topography; and any planting or removal of vegetation at a site that may be undertaken in association with these procedures (Group 70 International 2000). The 2000 Master Plan is somewhat vague on the definition of minor and major projects and leaves to the President of the University the final determination how the project would be classified. However, the operating definition considers construction activities – including excavation or the construction of new buildings – to be "major projects", while "minor projects" are those such as building small structures (e.g., weather tower) on a previously modified surface, or an emergency staircase. The 2000 Master Plan established separate review processes for minor and major projects. Minor project review ends with the UH President, while major projects require formal approval by the UH Board of Regents. OMKM functions as a liaison to ensure consistency in the project review process and that the required entities engage in the process. A schedule for processing proposals submitted to OMKM was approved in January 2008 (see Figure 5-1).

KKMC: Kahu Ku Mauna Council MKMB: Mauna Kea Management Board OMKM: Office of Mauna Kea Management Applicant submits proposal/request to OMKM Major Project MKMB Initial project classification and approval/disapproval 15 DAYS AFTER Recommendation for Major RECEIPT OF Project classification and PROPOSAL/REQUEST Preparation of an EA/ EIS is submitted to UHH Chancellor UH President Minimal Impact or Minor Project and determine whether project is exempt or requires an EA Disagrees Agrees OMKM initiates Major Project review process. KKMC reviews (Length of process varies according to complexity of project) Submits recommendation/comments to OMKM 30 Days after receiving proposal/request from OMKM OMKM reviews input from KKMC Submits recommendation to MKMB seven (7) days prior to next MKMB Recommends reclassifying Disapproves Minimal Minor Project to Major Impact project. Project MKMB Classification denied Recommends Preparation of an EA Approve Minimal Impact (Follow same steps for an Recommends Minor project classification or reclassifying EA under Major Project Review) Minimal Impact to Minor Project. No EA Applicant allowed to proceed with project Recommendation regarding Minor Project classification and project and project approval/disapproval submitted to UHH Chancellor, UH President (final approval) 10 days following MKMB meeting Approved by the OMKM notifies Applicant of Mauna Kea Management Board Decision on January 29, 2008

Figure 5-1. Schedule for Processing Proposals Submitted to OMKM




Figure 5-2. Major Project Review Steps

In January 2008, the MKMB approved "Major Project Design Review Steps" (see Figure 5-2). This flow chart details the Master Plan Design Review Process, EA/EIS Process, Master Plan Project Approval Process, and DLNR CDUA Process. Included in the overall design review process is the need to conduct required environmental analyses, such as an environmental assessment or environmental impact statement, for future proposed actions. As an advisory body to the MKMB, the Environment Committee plays a role in the project review process, especially with evaluating the potential impacts on natural resources.

5.1.2 Personnel, Training, Equipment, and Facilities

5.1.2.1 Staffing Requirements

Table 5-1 details the staffing requirements for oversight of the NRM Program and implementation of the NRMP. Most of the positions are existing, funded OMKM positions or volunteer boards or committees. A minimum of two new full-time OMKM positions are recommended for developing the NRM Program: a Natural Resources Coordinator and at least one field biologist. The number of field biologists needed will depend in part on what proportion of the natural resources management actions is conducted in house by OMKM and what proportion is conducted by paid contractors (i.e., subject matter specialists, environmental consultants, Bishop Museum scientists, University of Hawai'i faculty, and graduate students) and volunteer groups. As of the first draft of this NRMP, there are no dedicated natural resources management personnel within OMKM—all natural resources management activities are currently conducted by contractors advised by the MKMB Environment Committee.

Position Role **Current Status** Location **Funding Status OMKM Director** Oversight, guidance Exists and filled **OMKM Hilo** Currently funded (Interim director) **OMKM** Associate Oversight, guidance Exists and vacant OMKM Hilo Currently funded Director Mauna Kea Oversight, guidance Exists and filled⁵ Various (meets Unfunded at OMKM Hilo) Management Board volunteers MKMB Environmental Oversight, guidance Exists and filled⁵ Various (meets Unfunded Committee at OMKM Hilo) volunteers Kahu Kū Mauna Cultural guidance on Exists and filled⁵ Various (meets Unfunded NRMP issues at OMKM Hilo) volunteers Natural Resources Implementation of Position does not yet OMKM Hilo/VIS Requires funding NRMP Coordinator Natural resources field Implementation of Position does not yet OMKM Hilo/VIS Requires funding biologist(s) **NRMP** exist Natural resources Implementation of Needed Various Unfunded volunteers NRMP⁶ volunteers, but requires oversight by NRC, field biologist, or rangers.

Table 5-1. Staff Requirements

⁻

⁴ It is recommended that hydrological and geological studies be conducted by subject matter experts under contract.

⁵ See Section 1.4.2.1 for more information on the MKMB and its associated committees, OMKM staff (director, administrators, librarian), rangers, Mauna Kea Support Services (MKSS), and Visitor Information Station (VIS) staff.

⁶ Management actions conducted by volunteers could include invasive plant removal (weeding), establishing native gardens and maintaining them, education and outreach (on the mountain and at schools), plant and animal surveys (by qualified individuals), trail maintenance, and trash and debris removal. Activities that occur in the field would require a safety orientation and supervision by OMKM natural resources staff or rangers, to ensure safety of volunteers.

Position	Role	Current Status	Location	Funding Status
Contractors (scientists, environmental consultants, GIS database management)	Implementation of NRMP ⁷	Needed ⁸	Various	Requires funding (project specific)
Rangers	Visitor safety and outreach; natural resources protection	Exists and filled ⁵	VIS	Currently funded
VIS Staff	Visitor outreach and education	Exists and is filled through MKSS ⁵	VIS	Currently funded staff and unfunded volunteers
Education and outreach coordinator	Visitor outreach and education	Position does not yet exist ⁹	OMKM Hilo, VIS	Requires funding
Administrative staff	Administrative duties	Exists and filled ⁵	OMKM Hilo	Currently funded
Librarian	Document retrieval and organization	Exits and filled ⁵	OMKM Hilo	Currently funded
MKSS	Facility support	Exists and filled ⁵	Hale Pōhaku, VIS	Currently funded

5.1.2.2 Training Requirements

Training requirements for all OMKM personnel involved in field-based natural resource management activities includes general safety training, 4-wheel drive vehicle operation, orientation to working at high elevations, CPR and first aid, Global Positioning System (GPS) operation, rules and regulations, and recognition of culturally significant areas and items and protected flora and fauna. Additional training or educational needs for specific personnel are listed below.

Natural Resources Coordinator: Prior to hiring, the NRC should have extensive training in natural resources management issues, preferably in the form of a graduate degree and relevant work experience. On hiring, initial training would include training in project management, education and outreach methods appropriate to Hawai'i, and familiarization with the plant and animal species found on Mauna Kea, with emphasis on protected species. Other activities to maintain current working knowledge of natural resources management issues would include regular attendance of scientific meetings and conferences, participation in local working groups, and review of relevant scientific journals. Computer training for the NRC should include, if needed, use of statistical packages, Geographic Information System (GIS) software, word processing and database software. Optional training: GIS database management, spatial analysis, and advanced statistics.

Field biologists: Field biologists should have, at minimum, an undergraduate degree in biology or a related field. They should be familiar with field techniques to be used in day-to-day natural resources management activities and should be trained to recognize common Mauna Kea plant and animal species, as well as protected species. Optional training: GIS database management, relevant computer software.

⁷ A range of management actions could be conducted by contractors, including GIS services, baseline inventory, hydrological studies, biological studies, various management actions, or research projects.

⁸ There are currently several natural resource related contracts underway, including studies of the status of the wēkiu bug by Bishop Museum scientists, and a study of wēkiu bug population dynamics, genetics, and natural history by a University of Hawai'i professor and graduate student. See Section 2.2.2.2 for more information on these studies.

⁹ New position recommended to coordinate non-astronomy related education and outreach activities.

Volunteers: Safety orientation; natural resources orientation (to be conducted by NRC or a field biologist).

VIS Staff and volunteers: Safety orientation; natural resources orientation (to be conducted by NRC or a field biologist).

5.1.2.3 Facilities and Equipment

Facilities and basic equipment needed to implement the NRMP include

Facilities Equipment

- Office space for NRC and field biologist at OMKM, Hilo
- Equipment storage area at Hale Pōhaku
- Meeting room with audio-visual equipment at OMKM Hilo
- Library space for NRMP-related books and documents at OMKM Hilo
- Computers for NRC and field biologist, with database software, word processing, and GIS software
- Color printer
- Digital cameras (2 for natural resources staff, 1 for ranger truck)
- Walkie-talkies (minimum of 4 sets)
- GPS Units (2 for natural resources staff, 1 for ranger truck)
- 4-wheel-drive truck
- Field equipment (backpacks, measuring tapes, metal stakes, flagging, compasses, field notebooks, binoculars, sample collection materials)
- Identification guides (plants, birds, arthropods)

5.1.3 Ongoing Coordination with Other Agencies

Coordination between state and federal agencies regarding management of Mauna Kea's natural resources must be improved. This section describes two programs for coordinating multi-agency management activities on the mountain. The first (Section 5.1.3.1) is an interagency working group to streamline management activities on the mountain and to ensure that management occurs at an ecosystem level. The second (Section 5.1.3.2) is an annual interagency meeting to review and comment on the success of the OMKM NRM Program and the NRMP. Finally, Section 5.1.3.3 describes additional interagency coordination efforts that should be continued or undertaken to ensure the success of the NRM Program.

5.1.3.1 Mountain-wide Natural Resources Management Coordination

The principles of ecosystem management require that neighboring landowners and managers work together, guided by well-established management goals and visions. Overlapping and adjacent jurisdictions at the high elevations of Mauna Kea involve multiple agencies in management and decision-making (see Section 1.4.2). A good example of interagency coordination for land management on the Island of Hawai'i is the Three Mountain Alliance (TMA). It is recommended that the TMA management plan be reviewed and that prior to establishing the Mauna Kea working group, advice be sought from key personnel involved in the formation and operation of the TMA (Three Mountain Alliance 2007).

To increase participation and collaboration between OMKM and state and federal agencies on Mauna Kea, the following actions are recommended

- 1. Participate in the development of an interagency working group involving all entities that are responsible for or involved in natural resources management in high elevation areas (above 6,200 ft, or 1,900 m)¹⁰ on Mauna Kea, including OMKM, state and federal agencies, non-governmental organizations (NGOs), private land owners, and other agencies and persons involved in the day-to-day management of Mauna Kea lands.
 - Table 5-2 lists potential participants in the working group. If such a group already exists when the NRM Program is established, it is recommended that OMKM join it, and the NRC attend meetings.
- 2. Hold annual working group meetings.
 - a. During the first annual meeting
 - i. Develop an interagency set of mountain-wide management goals based on the principles of ecosystem management. Goals will need to take into account the participants' differing approaches to resource management, their policy foundations, and the decision criteria used by different institutions involved in multi-agency planning processes. Factors that will have to be taken into account include
 - 1. Department of Land and Natural Resources (DLNR) rules and regulations, including those specific to
 - a. Natural Area Reserve (NAR)¹¹
 - b. Mauna Kea Forest Reserve
 - c. Lands leased to UH¹²
 - d. Conservation District lands
 - e. Conservation District Use Permit regulations and stipulations
 - 2. Department of Hawaiian Home Lands rules and regulations
 - 3. Department of Defense, U.S. Army, Pōhakuloa Training Area (PTA) rules and regulations
 - 4. U.S. Fish and Wildlife Service regulations regarding protected species
 - ii. Clarify management roles on the mountain: Who will carry out what activities and where the funding will come from. Managed activities could include
 - 1. Ecosystem restoration
 - 2. Protection and enhancement of protected species (e.g., silversword outplanting)
 - 3. Hunting
 - 4. Invasive species management

.

¹⁰ For simplicity's sake, the working group, at least in the beginning, should focus on the high elevation areas. The cut-off (6,200 ft) was chosen because it is recognized as the boundary above which subalpine vegetation begins on Mauna Kea (Mueller-Dombois and Fosberg 1998). However, the working group may decide to change the area covered once it convenes.

¹¹ OMKM has developed a Cooperative Agreement to address cross-boundary management issues related to the Mauna Kea Ice Age NAR (see Section 1.3.3.1).

¹² As part of the lease agreement, DLNR and UH share responsibility for management in the Mauna Kea Science Reserve (MKSR) and at Hale Pōhaku. See Table VIII-1 in the 2000 Master Plan (Group 70 International 2000) for a detailed matrix showing responsibilities in the high elevation areas of Mauna Kea. These responsibilities may change if UH is able to obtain rule-making authority.

- 5. Access
- 6. Recreation
- 7. Trail and road maintenance
- 8. Commercial enterprises (e.g., sightseeing tours)
- iii. Identify opportunities for collaboration, cooperation, and streamlining of natural resource management activities on the mountain.
- iv. Identify gaps in management activities and funding, and collaborate in efforts to fill the gaps.
- v. Identify time frames for various management projects and activities.
- b. At subsequent annual meetings
 - i. Discuss progress towards identified goals.
 - ii. Determine causes of any delays in meeting goals and any problems with implementation.
 - iii. Revise goals, management roles, time frames, and methodology, as needed.
 - iv. Identify new sources of funding, funding gaps, and new potential partners.

Table 5-2. Potential Participants in Mauna Kea Natural Resources Management Working Group

Source	Agency
Federal: Landholders on Mauna Kea ¹³	U.S. Army, Pōhakuloa Training Area (PTA)
Federal: Advisory or Regulatory	U.S. Fish and Wildlife Service (USFWS)
Agencies	U.S. Geological Survey (USGS)
	U.S.D.A. Forest Service, Institute of Pacific Islands Forestry (IPIF)
	U.S.D.A. Natural Resources Conservation Service (NRCS)
State: Landholders/leaseholders	University of Hawai'i (OMKM, IfA)
	DLNR: Division of Fish and Wildlife (DOFAW) and Natural Area Reserves System (NARS)
	Department of Hawaiian Home Lands (DHHL)
State: Advisory or Regulatory	Big Island Invasive Species Committee
Agencies	DLNR: DOFAW
	BLNR
	Office of Hawaiian Affairs
NGO	Silversword Alliance
	The Nature Conservancy
	Three Mountain Alliance
	Sierra Club
	Hawaii Watershed Alliance
	Hawaii Conservation Alliance
Individuals	Local natural resource experts
	Private landowners
	Community members
	Conservation advocates

 $^{^{\}rm 13}$ Landholders above 6,200 ft (1,900 m) elevation.

-

5.1.3.2 OMKM Natural Resources Management Program Interagency Review

OMKM personnel (including OMKM director, NRC, and members of the MKMB Environment Committee) should meet annually with relevant state and federal agencies to review the status of the OMKM NRM Program. Before the meeting, the NRC should prepare an annual progress report that describes the state of the resources, the status of the management program, progress towards meeting annual goals, and other pertinent information. The report will be reviewed and approved by the MKMB Environmental Committee and OMKM and will then be submitted to the stakeholders and agencies participating in the review process, allowing ample time before the meeting for the agencies to review it. The annual meeting will provide a mechanism for various agencies to review the OMKM NRM Program, provide feedback on management activities, and suggest additional activities or changes to the management program. Agencies to involve include USFWS, DLNR, and any other state or federal agencies participating in management activities or agreements with OMKM. The annual progress report is discussed further in Section 5.2.

5.1.3.3 Other Interagency Efforts

Other interagency efforts that will increase the success of the NRM Program include

- 1. Continue participation of the Big Island NARS manager on the MKMB Environment Committee.
- 2. Ensure diverse representation on the MKMB Environmental Committee, including local scientific experts, conservation advocates, and land managers and agencies.
- 3. Establish working relations with environmental staff at PTA, to cooperate and coordinate on natural resources management projects.

5.1.4 Budgets and Funding

Start-up costs for the NRM Program include furnishing office space and storage areas and for purchasing basic equipment (see Section 5.1.2.3). Annual operational expenses for the NRM Program will include salary and benefits for the NRC and field biologist(s), maintenance of equipment and vehicles, expenditures for various contracts¹⁴ (initial baseline inventory, specialized annual monitoring¹⁵, GIS database management). Annual operational expenses should also include a travel allowance for the NRC to attend public meetings on-island and interisland, the Hawai'i Conservation Conference, and one other relevant conference annually. Although many of the management actions can be implemented by operational funds to support staff, project-specific funds may be required for one-time projects or projects that require large amounts of funding, such as baseline inventories, mitigation or large-scale restoration projects, and multi-agency projects.

Potential sources of funds for start-up and annual operation costs include

- Increase in base funding provided to OMKM by the University of Hawai'i
- OMKM Revolving Fund, which is supported in part by fees collected from commercial tour operators

_

¹⁴ If not done in-house.

 $^{^{15}}$ Specialized annual monitoring is monitoring that requires in-depth knowledge that the NRM may not possess, such as identification of certain organisms or plants.

- Funds provided by existing observatories as good-will efforts, to aid natural resources management of the summit and Hale Pōhaku
- Funds provided through lease agreements for potential new astronomy development to support natural resources management activities
- User fees and licenses: vehicle entry fees (if implemented); license fees from commercial tour operations; fees for rental or use of facilities at Hale Pōhaku for non-astronomy activities, events, and research
- Revenue collected from enforcement of rules (if implemented)

Potential sources of project-specific funding include

- OMKM base-funding requests
- Cost-sharing through agency partnering
- Using volunteer labor when possible
- Funds provided by new projects and existing observatories as part of environmental analysis or mitigation of environmental damage
- Research and management grants provided by federal and state agencies

5.1.5 Schedule

Implementation of the NRMP will begin with the acceptance of the NRMP by the MKMB and BLNR (as part of the Mauna Kea Comprehensive Management Plan [CMP]). The development of the NRM Program would begin once the NRMP is accepted and funding sources are identified. ¹⁶ The following schedule begins once funding is secured for the NRM Program, including funding for baseline inventories.

Year 1

- 1. Obtain office space and equipment needed for NRM Program (see Section 5.1.2.3).
- 2. Advertise and fill NRC and field biologist positions (see Sections 5.1.2.1 and 5.1.2.2).
- 3. Request proposals for baseline inventories (plants, invertebrates, birds and mammals, hydrological and geological resources) and hire contractors.
- 4. Begin development of NRM Program.
 - a. Begin developing component programs as described in the Section 4 component plans
 - b. Begin education and outreach activities
- 5. Draft RFPs and determine funding sources for projects that cannot be completed in house.

Year 2

- 1. Complete baseline inventories and associated data analysis.
 - a. Using data obtained from baseline inventories, designate high-priority management areas (e.g., areas with high ecological value, sensitive areas, critical habitat or rare species habitat,

 $^{^{16}}$ OMKM may need to secure increases in base funding or obtain funds from other sources. If funding is limited, this NRMP may be implemented incrementally.

and areas with significant ecological damage or problems). These areas will be the main focus of management activities.

- 2. Determine high-priority management actions for each of the component plans, using information obtained from baseline inventories.
- 3. Continue developing in-house programs, such as education and outreach activities, invasive species prevention and control.
- 4. Establish volunteer groups and conduct volunteer-based resource management activities.

Year 3

- 1. Review, evaluate, and revise NRMP as needed, to reflect findings from baseline inventories.
- 2. Establish priorities for management actions and develop a long-term timeline for natural resources management projects.
- 3. Begin annual monitoring (see Section 4.1).
- 4. Continue and improve upon component programs.

Year 4

- 1. Review, evaluate, and revise NRM Program as needed, based on Year 3.
- 2. Continue and improve upon component programs.

Year 5

- 1. Continue and improve upon component programs.
- 2. Review and evaluate NRM Program, in-depth.
- 3. Prepare stakeholder report on the status of Mauna Kea's natural resources, and the progress of the NRM Program.

Year 6

- 1. Continue and improve upon component programs.
- 2. Hold NRMP review meetings (see Section 5.2.2.2).
- 3. Revise NRMP, reprioritize NRM Program actions for following five-year interval.

5.2 Evaluation Plan

To ensure the best possible protection for Mauna Kea's high-elevation resources, the success of management actions to protect and conserve those resources must be regularly evaluated. This section describes methods that can be used to ensure that management actions are achieving stated NRMP goals and objectives in UH Management Areas. Section 5.2.1 describes how the NRC and OMKM will monitor progress towards meeting natural resources management goals. Section 5.2.2 describes the NRMP review and revision process.

5.2.1 Monitoring NRMP Implementation

Regular monitoring of NRMP implementation must occur in order to determine if progress is being made towards meeting the goals and objectives of the NRMP. The NRM Program will be monitored annually,

with annual progress reports issued as described in Section 5.2.2.1. A major review and revision of the NRMP will occur every five years, as described in Section 5.2.2.2.

Monitoring of a natural resources management program requires collection of concrete data that can be objectively analyzed and compared between years. This requires preparation at the beginning of each year, to determine the performance measures by which the year will be judged, followed by collection of data throughout the year on progress made toward meeting these measures. Preferably, the NRC will conduct brief monthly progress checks, to ensure that management activities are begun at appropriate times, that nothing is forgotten, and that projects are progressing as scheduled. Good notes taken throughout the year on the causes of delays or concerning unrealistic scheduling will enable the NRC to more thoughtfully analyze annual progress and determine course-corrections for the following year. The annual monitoring program should occur as follows:

- 1. At the beginning of each year, the NRC will establish a list of priority management actions to occur that year, along with a realistic schedule. These actions should be based on the components developed in the NRMP, along with any new pressing issues that may have come to light since the last update of the NRMP.
 - a. The NRC should take care that the schedule is realistic and that there are not more tasks than can be completed in the number of man-hours available in the year. If it is consistently found that there are more tasks that must be completed than there are man-hours to complete them, staffing needs should be reviewed and new staff added, as needed.
 - b. Management actions and projects that require a great deal of field labor (e.g., invasive weed control on a large scale), or expertise that the NRC and field biologist(s) do not have (e.g., invertebrate surveys if the NRC or field biologist is not an entomologist) should be contracted out, to ensure that they are conducted on schedule.
- When the list of priority management actions and the schedule is completed, the NRC should create a progress tracking datasheet, using the listed actions, and begin tracking progress towards meeting the actions.
 - a. It is useful to break up larger actions into smaller components that can be individually tracked and checked off when complete. This will give a sense of progress for some of the larger items that may be complete only at the end of the year, or even after several years. Ongoing projects should be broken into monthly components, or as appropriate.
 - b. As each action is completed, the NRC should enter the date of completion into the datasheet. It is easy to forget when projects were completed if you are attempting to recall this information at the end of a busy year.
 - c. Notes on problems encountered during management actions, interesting outcomes, successes, and ideas for improving management actions in the future should be kept on a linked document, to allow for easy cross-reference. This will help when writing the progress report at the end of the year.
 - d. The progress tracking datasheets and schedule should be referenced at the beginning of each month, and updated as appropriate.
- 3. At the end of each year, the NRC should review the progress of that year's management-program activities by reviewing the progress-tracking datasheet. This datasheet will provide information on the percentage of management actions completed during the year and can reveal patterns in the strengths and weaknesses in the management program. The notes will provide helpful information on how best to improve the management actions, if they are to be continued the next

year. Comparison of the projected schedule with the actual schedule will enable the NRC to better estimate timelines for future projects and will help determine if the scale and scope of the actions slated for the following year(s) are appropriate for the staffing level.

4. After the progress tracking forms are analyzed, the NRC should produce an annual report, as described in Section 5.2.2.1.

5.2.2 Review and Revision

The principles of adaptive management require regular review of the program and revision of management goals, objectives, actions, and techniques, to improve the performance of the program. Two review processes, an annual progress report and a five-year management outcome assessment, are recommended to assess the success of the NRM Program and to enable revision of the NRMP.

5.2.2.1 Annual Progress Report

At the end of each year the NRC should produce an annual progress report describing in detail the management goals, objectives, and actions for the year and what progress was made towards meeting them. The report should also describe actions to be taken to improve the program for the next year(s). This report is not intended to be a status report on the natural resources in UH Management Areas, ¹⁷ rather it is meant to inform management and stakeholders of the progress of the program and direction it is to take in the future.

This progress report should be presented at an annual meeting, to be held before the following year's management priorities and schedule are set. This will allow for input into the NRM Program by OMKM, MKMB, the MKMB Environment Committee, DLNR, and other agencies, as deemed appropriate (see Section 5.1.3.2).

5.2.2.2 Five-Year Management Outcome Analysis and NRMP Revision

The NRM Program should be subjected to a major review every five years, and the NRMP should be revised, as necessary. This process should involve input from state and federal agencies and the public.

5.2.2.2.1 Management Outcome Analysis

Determination of the outcome of management activities on the natural resources and of the success of the management program will be accomplished through a report summarizing the state of natural resources on Mauna Kea and the progress of the NRM Program over the preceding five years.

The first section of the report will discuss the state of the natural resources in UH Management Areas. This section will summarize data collected during monitoring, research, restoration, and threat prevention and control activities conducted over the preceding five years (See Section 4). This portion of the report will analyze trends in natural resources, and the impacts (positive, negative, or neutral) that management actions have had on them. It will also summarize what future management actions are needed to protect, enhance, or restore Mauna Kea's natural resources.

_

¹⁷ The status of natural resources in UH Management Areas will be tracked during annual natural resources monitoring and will be presented in the annual Natural Resources Monitoring Report. See Section 4.1 for information on natural resources monitoring.

¹⁸ The source of the data will be the annual reports produced as part of the various component plans.

The second section of the Five-Year Management Outcome Analysis should include a summary of the progress of the NRM Program towards meeting management goals, objectives, and actions, as outlined in the NRMP and in the annual listing of priority management actions (see 5.2.1). The source of information for this section of the report will be the annual progress reports from the last five years. Additionally, the NRC should review the NRMP and determine if any goals or objectives were not addressed during the preceding five years, and if so, why not. This section will discuss strengths and weaknesses of the NRMP and the NRM Program and ways to improve them.

The purpose of the Management Outcome Analysis is to provide analysis of both the condition of the natural resources in UH Management Areas and the status of the NRM Program and Plan. This information will be used to update the NRMP, so that it better addresses the needs of the natural resources, and to improve management activities through adaptive management.

A draft of this report should be submitted for review and comment to OMKM, MKMB, the MKMB Environment Committee, and state and federal agencies, as deemed necessary or appropriate. This will provide a mechanism for the interested parties to provide input into the direction the NRM Program and suggestions for changes to the NRMP. A final version of the report can then be presented to the public for comments and suggestions to be used in revising the NRMP.

5.2.2.2.2 Revising the Natural Resources Management Plan

Following the production of the Five-Year Management Outcome Analysis, and after input from appropriate stakeholders, the NRMP should be revised and updated to incorporate new management goals, objectives, and actions. This major review and revision process should occur on the sixth year (to allow for time to process the five-year review). If it is determined that the five-year cycle is too short to show real changes in resource conditions, then after two five-year review and revision cycles, the frequency of the process can be lengthened, as needed.

It should be noted that as of the first draft of this NRMP, no quantitative baseline inventories of the natural resources on UH Management Areas have been conducted. Therefore, the true state of the natural resources is currently unknown, and completion of the baseline inventories will necessitate a reevaluation of the management actions recommended in the component plans in Section 4. It may be necessary to complete one or more in-house reviews and revisions of the NRMP and management priorities during the first several years of the program, to determine impediments to successful management of natural resources, develop realistic timelines for projects, and make necessary changes to the structure of the program. This can be done on an as-needed basis, to be determined by the NRC, OMKM director, and MKMB Environment Committee.

6 List of Preparers

Ken Banks, Editor

Colleen Bergmanis, M.S., Natural Resource Management Specialist

Kristin Duin, M.S., Principal

Jennifer Garrison, PhD, Senior Natural Resource Management Specialist

Kehaupuaokalani Kamaka, Research Assistant

Wendy Kuntz, PhD, Senior Natural Resource Management Specialist

Andrew Hood, M.S., Principal

Michelle Roberts, M.S., Natural Resource Management Specialist

Krista Winger, Island Geographic, LLC, Geographic Information Systems

Section 6. List of Preparers

This page intentionally left blank.						

7 List of Persons Consulted

The following individuals were consulted during the development of this Natural Resources Management Plan, either through personal communication, interviews, surveys, or attendance at public meetings.

Teddy Akau, Kona Lua

Alison Ainsworth, DLNR NARS

Patrick Aldrich, Department of Zoology, University of Hawai'i

Shahin Ansari, SWCA Environmental Consultants, Honolulu

Greg Asner, Stanford University

Douglas Astrande, MKSS

Ron Bachman, DLNR DOFAW

Paul Banko, U.S. Geological Survey

John Barnes, Director, NOAA Mauna Loa Observatory

Randal Bartlett, Pu'u Kukui Watershed Preserve, Maui Land & Pineapple Company, Inc.

Steven Bergfeld, DLNR DOFAW

Paul Berkowitz, U.S. Geological Survey

Tom Birchard, National Weather Service Forecast Office

Frank Bonaccorso, U.S. Geological Survey

Gregory Brenner, Pacific Analytics LLC

Mike Britten, Rocky Mountain I&M Network, National Park Service

Jackie Brown, Grinnell College

James Bruch, Kaho'olawe Island Reserver Commission

Marie Bruegmann, U.S. Fish and Wildlife Service/Pacific Islands Ecosystem Conservation

Steve Businger, Department of Meteorology, University of Hawai'i

David Byrne, Manager, Mauna Kea Visitor Information Station

Curt Cottrell, DLNR State Parks

Steve Clark, Pacific Consulting Services, Inc.

Pat Conant, Hawai'i State Department of Agriculture

Sheila Conant, Department of Zoology, University of Hawai'i

Ian Cole, DLNR DOFAW/NARS

Susan Cordell, USDA Forest Service Institute of Pacific Islands Forestry, MKMB Environment Committee

Bob Covington, DLNR DOFAW

Sara Da Silva, University of Hawai'i

Bethany Ehlmann, Brown University

Julia Ehses, University of Hawai'i

Jesse Eiben, Plant and Environmental Protection Sciences, University of Hawai'i

Ron Englund, Kamehameha Schools

Steve Evans, Pohakuloa Training Area

Section 7. List of Persons Consulted

Steve Fancy, National Park Service, National Monitoring Program

Chris Farmer, U.S. Geological Survey

Lisa Ferentinos, DLNR DOFAW

Brad Finney, Environmental Resources Engineering, Humboldt State University

David Foote, U.S. Geological Survey

Shane Fox, OMKM Ranger

Scott Fretz, DLNR DOFAW

Betsy Gagné, DLNR NARS

Tom Giambelluca, Geography Department, University of Hawai'i

Sam Gon, The Nature Conservancy

Daniel Gruner, University of Maryland

Michael Hadfield, Department of Zoology, University of Hawai'i

Lisa Hadway, DLNR NARS, MKMB Environment Committee

Kevin Hamilton, International Pacific Research Center (IPRC), University of Hawai'i

Saeko Hayashi, Subaru Telescope

Leslie HaySmith, National Park Service Inventory & Monitoring Program, Volcano

Don Hemmes, Biology Department, University of Hawai'i at Hilo

Jennifer Higashino, DLNR DOFAW

Nelson Ho, Sierra Club; Department of Environmental Management, Hawai'i County

Frank Howarth, Bishop Museum

Roger Imoto, DLNR DOFAW

Jack Jeffrey, U.S. Fish and Wildlife Service

Ed Johnson, DLNR DOFAW

Irv Kawashima, DLNR DOFAW

Randy Kennedy, DLNR DOFAW

Susan Kihara, Hawai'i State Department of Health Clean Air Branch

Fritz Klasner, National Park Service Inventory and Monitoring Program, Volcano, Hawaii

Ron Koehler, Director, MKSS

Christoph Kueffer, Department of Botany, University of Hawai'i

Sigrun Kullik, Environmental Biology, University of Guelph

Tiara Lackey, Pōhakuloa Training Area

Sam Lemmo, DLNR OCCL

David Leonard, DLNR DOFAW

John Lockwood, Geohazards Consultants International, Inc., Volcano, Hawai'i

Rhonda Loh, Hawai'i Volcanoes National Park

Lloyd Loope, U.S. Geological Survey

Scot Loring, Pacific Crest Consulting, Talent, Oregon

Karl Magnacca, Department of Zoology, Trinity College, Dublin, Ireland

Kepa Maly, Kumu Pono Associates LLC

Christy Martin, CGAPS-Coordinating Group on Alien Pest Species

Wes Matsunaga, DLNR Land Division

Pat McCoy, Pacific Consulting Services, Inc.

Robert McLaren, Specialist and Associate Director, Institute for Astronomy

Matthew Medeiros, University of California, Berkeley

Joey Mello, DLNR DOFAW

Theresa Menard, The Nature Conservancy

Marty Meyer, Department of Zoology, University of Hawai'i

Stephen E. Miller, U.S. Fish and Wildlife Service

Christien Mitchell, DLNR DOFAW, NARS

Michelle Montgomery, U.S. Geological Survey

Kevin Moore, DLNR Land Division

Stephanie Nagata, OMKM

Ron Nahakuelua, MKSS

Miles Nakahara, DLNR DOFAW

Larry Nakayama, Planning Department, County of Hawai'i

Tom Nance, Tom Nance Water Resource Engineering

Jerry Nunogawa, Waste Water Branch, Hawai'i State Department of Health

Peter Oboyski, University of California, Berkeley

Dawn Pamarang, OMKM Secretary

Pablo McLoud, OMKM Ranger

Dan Polhemus, DLNR DAR

Thane Pratt, U.S. Geological Survey

David Preston, Bishop Museum

Rob Robichaux, University of Arizona

Jason Rohr, Department of Biology, University of South Florida

Joby Rohrer, Natural Resource Management Coordinator, U.S. Army

Rebecca Rundell, Botany Department, University of British Columbia

Hans Sin, DLNR DOFAW

Clifford Smith, University of Hawai'i Manoa Emeritus

Fred Stone, University of Hawai'i Hilo Retired

Bruce A. Snyder, Odum School of Ecology, University of Georgia

Forest and Kim Starr, U.S. Geological Survey/RCUH

Andrew Taylor, Department of Zooogy, University of Hawai'i

Ron Terry, Geometrician Associates, MKMB Environment Committee

Don Thomas, Hawai'i Institute of Geophysics and Planetology, University of Hawai'i, MKMB Environment Committee

James Toma, Indoor and Radiological Health Branch, Hawai'i State Department of Health

Andrew Tomlinson, Pacific Consulting Services, Inc.

Eric VanderWerf, Pacific Rim Conservation

Visitor Information Station Staff

Section 7. List of Persons Consulted

Lorena Wada, U.S. Fish and Wildlife Service
Debbie Ward, Sierra Club, *MKMB Environment Committee*Richard Wass, Hakalau Forest NWR
Lauren Weisenberger, Botany Department, University of Hawai'i
Susan Will-Wolf, Department of Botany, University of Wisconsin
Kealoha Wilson, Logistics Clerk, MKSS
Joan Yoshioka, PEP Program, DLNR
Lindsay Young, Department of Zoology, Univesity of Hawai'i
Marjorie Ziegler, Conservation Council for Hawai'i

8 Glossary

Aa: Common type of basalt lava with rough broken surface; from Hawaiian word 'a'ā.

Adaptive management: Systematic process for continually improving management policies and practices by learning from the outcomes of past and current management activities. Adaptive management recognizes that there is a level of uncertainty about the "best" policy or practice for a particular management issue, and therefore requires that each management decision be revisited in the future to determine if it is providing the desired outcome.

Aeolian: Process pertaining to the wind in which particles are detached, transported and deposited. On Mauna Kea the aeolian processes are integral to at least one endemic arthropod due to the import of food resources transported from lower slopes of Mauna Kea to its highest elevations.

Albedo: Ratio of the amount of sunlight that is reflected and absorbed by a surface; 1.0 is a completely reflective surface.

Alien: When used in reference to a plant or animal species, alien means that the species was introduced (accidentally or on purpose) by mankind, and does not naturally occur in the location.

Alkalic: Igneous rock chemical classification based on total alkali (Na₂O+K₂O) vs. silca (SiO₂) content; alkalic rocks have greater total alkali than tholeitic rocks at the same silica content.

Alpine: Living or growing on mountains above the timberline. On Mauna Kea alpine ecosystems occur above the nocturnal ground frost line, which is found at 9,800 ft (3,000 m).

Alteration: Change in character, appearance, direction, or status.

Anthropogenic: Relating to or resulting from the influence humans have on the natural world.

Aquifer: Water bearing geological layer. Aquifers are often used to provide water for human needs.

Arthropods: Organisms with jointed appendages and exoskeletons, including insects, spiders, and crustaceans.

Baseline inventory: Initial survey establishing the status of natural resources (population sizes, distributions, and biological diversity). Conducted at the beginning of a natural resources management program.

Bedload: Particles of sand, gravel, or soil moved by water along the bottom of a stream, gulch or river.

Biodiversity: Full range of natural variety and variability within and among living organisms, and the ecological and environmental complexes in which they occur.

Candidate species: Species that is possibly declining in population and that is being considered for threatened or endangered status under the Endangered Species Act.

Cesspool: Underground reservoir for liquid waste.

Cinder: Fragment of cooled pyroclastic lava material, typically very porous and with low particle density, making them light for their size. Also called tephra and scoria.

Cinder cone: Conical hill formed by the accumulation of volcanic ash or cinder around a vent.

Climate: Long term average of meteorological variables representative of the geographic area where the variables have been collected.

Community: In ecology, a community is an assemblage of populations of different species, interacting with one another.

Control: When used in reference to invasive species, control is the long-term reduction of invasive species density and abundance to an acceptable threshold.

Decommission: To remove from service.

Dike: Mass of intrusive igneous rock that has been injected into a fissure while molten, usually vertically aligned to the ground surface.

Dormant: Marked by a suspension of activity; temporarily devoid of identifiable activity.

Ecosystem: Dynamic system of living organisms (plants, animals, and microorganisms) within an area, the environment sustaining them, and their interactions. An ecosystem can range in size from a tiny site containing only a few species, such as an isolated lake, to a huge area containing thousands of species, such as a tropical rainforest.

Ecosystem health: Condition of an ecosystem in which its dynamic attributes are expressed within the range of activity normal for its stage of development.

Ecosystem integrity: Achieved when an ecosystem has the biodiversity characteristics of a reference ecosystem such as species composition and community structure, and is fully capable of sustaining normal ecosystem functioning.

Ecosystem restoration: The process of assisting the recovery of an ecosystem that has been degraded, damaged or destroyed.

Endemic: Native to, and restricted to, a particular geographical region. Highly endemic species are those with very restricted natural ranges; they are especially vulnerable to extinction if their natural habitat is eliminated or significantly disturbed.

Ephemeral: Description of stream or gulch hydrology in which surface water flow occurs only following periods of rainfall or snow melt.

Eradicate: To completely remove or destroy the entire population. When used in reference to invasive species, it refers to removal of the invasive species from an entire landmass, such as the entire Island of Hawai'i. Eradication is extremely rare in the case of invasive species, and generally occurs only on very small islands and in isolated locations, or with very small populations.

Erodibility: Potential of soil to be eroded or physically altered by the action of water or wind.

Erosion hazard: Rating or classification of the potential of a surface to erode; used to identify areas to avoid developing or take special precautions.

Erosivity: Potential of water and wind to detach and transport soil or rock.

Evapotranspiration: Loss of water from the combined effects of evaporation and transpiration from plants.

Evaporation: Phase change of water from liquid to vapor; classified as a loss in the hydrologic cycle.

Exclosure: Limited area from which animals such as sheep, pigs, or cattle are excluded by fencing. Fencing, and subsequent removal of non-native mammals, is a habitat management technique used in areas containing native plants or animals that are susceptible to grazing or predation by non-native mammals.

Exotic: Introduced from another part of the world; not native to the place where found.

Fauna: Animals or animal life of a region or environment.

Fissure: Narrow opening or crack of considerable length and depth, usually occurring from some breaking or forced parting.

Flora: Plants or plant life of a region or environment.

Geographic Information System (GIS): Computer software used to store, view, and analyze geospatial information. Outputs include analysis, maps and images.

Glacial member: Layers of glacial deposit used to describe the three glacial episodes that occurred on Mauna Kea.

Gully: Trench originally worn in the earth by running water and through which water often runs after rain.

Headcut: Location where a sudden change in ground elevation occurs, usually at the leading edge of a gully. Headcuts often result in rapid erosion and incision of the runoff channel.

Hydraulic conductivity: Quantification of the transmission rate of a liquid. Movement of water through soil or rock is a function of the materials conductivity.

Incidental take: Unknown or accidental killing or removing of an organism.

Incipient invasive species: Invasive species that has not yet become established. For this plan, refers to known invasive species found in Hawai'i that are not currently on UH Management Areas but that are thought to pose a threat to subalpine or alpine communities.

Indigenous: Native to a given region or ecosystem. An indigenous species differs from an endemic species in that it may occur in several locations in the world, while an endemic species is limited to one area or region.

Interstitial pores: Openings or voids contained in a soil or rock.

Invasive species: Non-native (alien) species whose introduction does or is likely to cause economic or environmental harm or harm to human health.

Introduced: When used in reference to a plant or animal species, the term means that the species was brought to a new location by man (either purposefully or accidentally).

In-situ: In the natural or original position or place.

Isotopic analysis: The analysis of the isotope composition of a sample.

Leachate: Solution or product that gradually moves through a matrix to a new location.

Leach field: The soil surrounding and immediately down gradient of a septic tank used to absorb leached septic liquids.

Lichen: Symbiotic associations of a fungus with a photosynthetic partner (either green alga or cyanobacterium) that can produce food for the lichen from sunlight. Usually found growing on rocks or trees.

Lithology: Character of a rock or rock formation as described by its components.

MKSR: Mauna Kea Science Reserve. An 11,288 ac (4,568 ha) parcel of state land above 11,500 ft (3,505 m) elevation on Mauna Kea, leased by the University of Hawai'i.

Mauna Kea Forest Reserve: Approximately 448,000 acre forest reserve and part of the State of Hawai'i Forest Reserve System. Circles Mauna Kea, and is found directly below the MKSR.

Metadata: Data that provides descriptive information about other data.

Mitigation: Elimination, reduction or control of the adverse environmental effects of a project. Mitigation includes restitution for any damage to the environment caused by such effects through replacement, restoration, compensation or any other means.

Monitoring: Recording changes to an entity over time in order to understand how actions, activities and ecosystem processes affect it. Monitoring is conducted as part of a plan to evaluate management prescriptions.

Natural Area Reserve (NAR): Part of DLNR, the reserves are created to preserve and protect representative samples of Hawaiian biological ecosystems and geological formations. The Ice Age Natural Area Reserve is located on Mauna Kea.

Native: Naturally occurring in a given area. When used in reference to plants and animals in Hawai'i, the term native means species that were not brought to the islands by mankind.

NRCS soil survey: Scientific document describing the physical and chemical attributes of soil.

Nunatak: Areas of cinder cones that remained ice-free by being above the surrounding glacier during the previous glaciation period.

Orographic: Associated with or induced by the presence of mountains, e.g., cloud lift and development.

Outcrop: To project from the surrounding soil.

Outplanting: Planting of greenhouse or pot-grown plants into an area.

Paleosol: Soil horizon from the geologic past; usually buried beneath other rocks or recent soil horizons.

Percolate: Movement of liquid through a permeable substance.

Permafrost: Layer of ice usually located beneath the ground surface, which is maintained year round.

Pu'u: Hawaiian name for hill. With respect to Mauna Kea, pu'u refers to the volcanic cinder cones found at the summit.

Rare: Occurring at low densities, or in limited locations.

Rehabilitation: Repair of habitat following a disturbance. Rehabilitation emphasizes the repair of ecosystem processes, productivity, and services, rather than reestablishment of historical plant and animal communities (which is the focus of restoration).

Relic: Something that has survived the passage of time.

Restoration: The actions or activities implemented in order to return damaged or altered habitat or a component of the ecosystem to a historical condition, including physical habitat attributes as well as community composition and native species abundance.

Rill: Small erosion feature created by water flow. With time rills can either join together to form a gully or expand into a gully.

Ruderal: Plants that colonize disturbed sites, bare soil, or waste piles.

Seep: Spot where water trickles out of the ground.

Segetal: Plants that grow inter-mixed with crop species.

Sensitive area: Area that is deemed in need of protection. Sensitive areas may include areas with a rare, Threatened, or Endangered species; a unique native community; or physical resources. Sensitive areas may also be areas prone to disturbance such as erosion or crushing of cinder.

Septic tank: Tank used in the collection of human waste and gray water in which solid materials settle to the bottom and liquid waste is discharged onto a leach field (an area lined with highly porous materials allowing effluent to seep into the ground).

Site recycling: Reuse of a previously developed site. In the case of the observatories, refers to removal of an old observatory and replacement with a new one.

Snowpack: Accumulation of naturally formed, packed snow over an area that usually melts during the warmer months.

Species of Concern: Those species that might be in need of conservation action, but that are not currently Listed or Candidate species. Species of Concern receive no legal protection and use of the term does not necessarily imply that a species will eventually be proposed for listing.

Spring: Ground water discharging out of the earth where the water table meets the ground surface.

Stratigraphy: Geology discipline dealing with the origin, composition, distribution, and succession of strata (layers of sedimentary rock); the arrangement of strata.

Subalpine: Of, relating to, or inhabiting high upland slopes and especially the zone just below the timberline. On Mauna Kea, the subalpine region occurs from approximately 5,600 ft to 9,800 ft (1,700 m to 3,000 m) elevation.

Sublimation: Process by which water in its solid phase is transformed directly to vapor phase without first passing through the liquid phase. Sublimation rates are highest in areas where relative humidity is low, dry winds are present, and at high elevations where atmospheric pressure is low and sunlight is strong.

Subsidence: To sink; to tend downward. Refers to the slow process of volcanic island sinking due to the continued accumulation of extruded material and associated weight.

Tephra: Fragments of volcanic rock and lava regardless of size that are blasted into the air by explosions or carried upward by hot gases in eruption columns or lava fountains. Tephra includes large dense blocks and bombs, and small light rock debris that have usually solidified prior to hitting the ground.

Tholeiitic: Igneous rock chemical classification based on total alkali (Na₂O+K₂O) vs. silica (SiO₂) content; tholeiitic rocks have less total alkali than alkalic rocks at the same silica content.

Threatened and Endangered Species: Plant or animal that is in danger of extinction throughout all, or a significant portion, of its range. Federally listed threatened and endangered species are those species that the US Fish and Wildlife Service has determined require federal action to protect them. Protection of Federally listed threatened and endangered species is mandated by the United States Endangered Species Act, 1973.

Ungulate: Hoofed, typically herbivorous, quadruped mammal.

Upwelling: Process or an instance of rising or appearing to rise to the surface.

Vascular plant: Plant with water and fluid conductive tissue (xylem and phloem); includes seed plants, ferns, and fern allies. Does not include fungus, lichens or mosses.

Weathering: Physical disintegration and chemical decomposition of earth materials at or near the earth's surface.

Section 8. Glossary

This page intentionally left blank.						

9 References

- Abouchami, W. and A. W. Hofmann. 2000. Long-term Pb isotope record of sources within the Hawaiian plume [abs.]. Eos, Transactions, American Geophysical Union supp. v. 81(no. 46): p. F1355.
- Agrawl, A. 1995. Dismantling the divide between indigenous and scientific knowledge. <u>Development and Change</u> **26**(3): 413-439.
- Aldrich, P. 2005. Biological information for Mauna Kea. Prepared for the Office of Mauna Kea Management by Patrick Aldrich, Zoology Department, University of Hawaii Honolulu, HI.
- Aldrich, P. 2007. Personal communication from Patrick Aldrich, University of Hawaii, to Jennifer Garrison, SRGII, regarding native arthropod populations on Mauna Kea.
- Alexander, W. D. 1892. Scientific expedition to Mauna Kea A week on the summit. <u>The Friend</u> 50(10): 74-75.
- Alstad, D. N., G. F. Edmunds, Jr., et al. 1982. Effects of air pollutants on insect populations. <u>Annual Review of Entomology</u> **27**: 369-384.
- Alt, J. C. 1993. Low-temperature alteration of basalts from the Hawaiian Arch, Leg 136. R. H. Wilkens, J. Firth and J. Bender. College Station, TX, Texas A&M University, Ocean Drilling Program. v. 136: p. 133-146.
- Amarasekare, P. 1993. Potential impact of mammalian nest predators on endemic forest birds of western Mauna Kea, Hawaii. <u>Conservation Biology</u> 7(2): 316-324.
- Anonymous. 1948. Hunting on the big island of Hawaii. Paradise of the Pacific 60(5): 26-27.
- Ansari, S. 2008. Personal communication from Shahin Ansari, SWCA Environmental Consultants, to Jennifer Garrison, SRGII, regarding presence of common mullein at MKSR.
- Ansari, S. and C. C. Daehler. 2000. Common mullein (Verbascum thapsus): a literature review Technical Report 127. Pacific Cooperative Studies Unit, University of Hawaii at Manoa Honolulu. Retrieved from http://www.botany.hawaii.edu/faculty/duffy/techrep.htm.
- Arvidson, R. E. 2002. Draft environmental assessment for the Outrigger Telescopes Project published by NASA in December 2000; Response to comments concerning the hydrology of Mauna Kea. <u>Outrigger</u>. St. Louis, MO, McDonnell Center for the Space Sciences, Dept. of Earth and Planetary Sciences, Washington University.
- Ashlock, P. D. and W. C. Gagne. 1983. A remarkable new micropterous Nysius species from the aeolian zone of Mauna Kea, Hawaii Island (Hemiptera: Heteroptera: Lygaeidae). <u>International Journal of Entomology</u> 25(1): 47-55.
- Asquith, A. 1995. Alien Species and the Extinction Crisis of Hawaii's Invertebrates. <u>Endangered Species UPDATE</u> 12(6).
- Atkinson, C. T., K. L. Woods, R. J. Dusek, L. Sileo and W. M. Iko. 1995. Wildlife disease and conservation in Hawaii: Pathogenicity of avian malaria (Plasmodium relictum) in experimentally infected Tiwi (*Vestiaria coccinea*). Parasitology 111: 859-869.
- Baker, B. B. and R. K. Moseley. 2007. Advancing treeline and retreating glaciers: Implications for conservation in Yunnan, PR china. <u>Arctic Antarctic and Alpine Research</u> 39(2): 200-209.
- Baker, M. B., S. Alves, et al. 1994. Microscopic petrography of the Hawaii Scientific Drilling Project core: initial results [abs.]. <u>Eos, Transactions, American Geophysical Union supp.</u> v. 75(no. 44): p. 707-708
- Baker, M. B., S. Alves, et al. 1996. Petrography and petrology of the Hawaii Scientific Drilling Project lavas: inferences from olivine phenocryst abundances and compositions. <u>Journal of Geophysical</u> Research **101**(B5): 11,715-11,727.
- Balakrishinan, N. and D. Mueller-Dombois. 1983. Nutrient studies in relation to habitat types and canopy dieback in the montane rain forest ecosystem, island of Hawaii. <u>Pacific Science</u> **37**(4): 339-359.

- Baldwin, E. D. 1890. A trip to the summit of Mauna Kea. <u>Hawaiian Annual for 1890</u>. T. G. Thrum. Honolulu, HI: p.54-58.
- Baldwin, E. D. 1915. A trip to the summit of Mauna Kea. Paradise of the Pacific 28(9): 17-19.
- Banko, P. C. 2006. Palila Restoration: Lessons from Long-term Research. USGS FS 2006-3104. USGS Pacific Island Ecosystems Research Center.
- Banko, P. C., C. Farmer, K. W. Brinck, J. Castner, J. Crummer, R. Danner, B. Frederick, B. Hsu, B. Muffler, S. Nash, D. Nelson, D. Pollock, M. Schwartzfeld, R. Stephens and K. Rapozo. 2005. Palila Restoration Project, 2005. Summary of Results, 1996-2005. United States Geological Survey Pacific Island Ecosystem Research Center, Kilauea Field Station Hawaii National Park, HI.
- Banko, P. C., J. M. Black and W. E. Banko. 1999. Hawaiian goose (Nene) (Branta sandvicensis). <u>The Birds of North America No. 434</u>. A. Poole and F. Gill. Philadelphia, PA, The Birds of North America Inc.
- Banko, P. C., P. T. Oboyski, J. W. Slotterback, S. J. Dougill, D. M. Goltz, L. Johnson, M. E. Laut and T. C. Murray. 2002. Availability of food resources, distribution of invasive species, and conservation of a Hawaiian bird along a gradient of elevation. Journal of Biogeography 29(5-6): 789-808.
- Banks, R. C., C. Cicero, J. L. Dunn, A. W. Kratter, P. C. Rasmussen, J. V. Remsen, Jr., J. D. Rising and D. F. Stotz. 2002. Forty-third supplement to the American Ornithologists' Union Checklist of North American Birds. Auk 119(3): 897-906.
- Barnes, J. E. 2008. Personal communication from J. Barnes, Director, NOAA Mauna Loa Observatory regarding collection of air quality information Colleen Bergmanis SRGII.
- Barrows, E. M. 1980. Robbing of exotic plants by introduced carpenter and honey bees in Hawaii, with comparative notes. Biotropica 12(1): 23-29.
- Bartram, E. B. 1952. <u>High altitude mosses from Mauna Kea, island of Hawaii</u>. Honolulu, Bishop museum.
- Bayman, J. M. and J. J. M. Nakamura. 2001. Craft specialization and adze production on Hawaii Island. Journal of Field Archaeology **28**(3/4): 239-252.
- Bayman, J. M., Moniz-Nakamura, Jadelyn J., Rieth, Timothy M., and Paraso, C. Kanani. 2004. Stone adze production and resource extraction at Pohakuloa, Hawaii Island. <u>Hawaiian Archaeology</u> **9**: 83-104.
- Beeson, M. H., D. A. Clague, et al. 1996. Origin and depositional environment of clastic deposits in the Hilo Drill Hole, Hawaii. Journal of Geophysical Research. v. 101: p. 11,617-11,629.
- Benning, T. L., D. LaPointe, C. T. Atkinson and P. M. Vitousek. 2002. Interactions of climate change with biological invasions and land use in the Hawaiian Islands: modeling the fate of endemic birds using a geographic information system. <u>PNAS (Proceedings of the National Academy of Sciences)</u> 99(22): 14246-14249.
- Benton, N. 2006. Fact Sheet: Fountain Grass. Retrieved Sept. 23, 2007, from http://www.nps.gov/plants/alien/fact/pese1.htm.
- Berger, A. J. 1972. *Hawaiian birdlife*. Honolulu, University of Hawaii Press.
- Berger, A. J. 1975. The Hawaiian Honeycreepers, 1778-1974. Elepaio 35(10): 110-118.
- Bergfeld, S. 2008. Personal communication from Steven Bergfeld, DOFAW, to Jennifer Garrison, SRGII, regarding Mauna Kea silversword restoration and management efforts.
- Berkes, F. 2008. Sacred Ecology. New York, Routledge.
- Billings, W. D. 1979. High mountain ecosystems: evolution, structure, operation and maintenance. <u>High altitude geoecology</u>, AAAS selected symposium: 12. P. J. Webber. Boulder, Colo., Published for the American Association for the Advancement of Science by Westview Press: 97-125.
- Birchard, T. 2008. Personal communication from T. Birchard, lead forecast/meteorologist of the National Weather Service Forecast Office Honolulu based at UH Manoa regarding climate regime at Mauna Kea. C. B. SRGII.

- Bishop Museum. 2007a. Mauna Kea Online: Species found at (elev = 9400 ft.) Hawaii. Hale Pōhaku; Timberline; Mauna Kea. Retrieved Sept. 17, 2007, from http://www2.bishopmuseum.org/mkmk2/natsci.asp?Locality=Mauna+Kea&Collection=Botany&offset = 50.
- Bishop Museum. 2007b. Mauna Kea Online: Species found at Hale Pohaku. Retrieved Sept. 17, 2007, from
 - http://www2.bishopmuseum.org/mkmk2/natsci.asp?Locality=Hale%20Pohaku&Collection=Botany.
- Bishop Museum. 2007c. Mauna Kea Online: Species found at Mauna Kea (Entomology Collection). Retrieved January 27, 2007, from http://www2.bishopmuseum.org/mkmk2/natsci.asp?Locality=Mauna%20Kea&Collection=Entomolog y.
- Bjork, R. G. and U. Molau. 2007. Ecology of alpine snowbeds and the impact of global change. <u>Arctic Antarctic and Alpine Research</u> 39(1): 34-43.
- Blanchard, D. C. 1953. Raindrop size-distribution in Hawaiian rains. <u>Journal of Meteorology</u> **10**: 457-473.
- Bonaccorso, F. 2008. Personal communication from Frank Bonaccorso, USGS, to Jennifer Garrison, SRGII, regarding Hawaiian hoary bat biology and monitoring techniques.
- Brady, N. C. and R. R. Weil. 2000. <u>Elements of the Nature and Properties of Soils</u>. New Jersey, Prentice Hall.
- Brenner, G. J. 2002. Wekiu Bug Monitoring Project. from http://www.malamamaunakea.org/environment.php?article_id=20.
- Brenner, G. J. and J. Lockwood. 2005. Wekiu bug habitat quantitative cinder evaluation, Prepared for the Outrigger Telescopes Project by Pacific Analytics LLC and Geohazards Consultants International: 44 p.
- Brenner, G. J., D. J. Preston and F. G. Howarth. 1999. An arthropod assessment within selected areas of the Mauna Kea Science Reserve. Final Report. Documentation of methodology. Contribution No. 1999-007 to the Hawaii Biological Survey. Prepared for the University of Hawaii Institute of Astronomy Honolulu, HI.
- Brenner, G. J., P. T. Oboyski and P. C. Banko. 2002. Parasitism of Cydia spp. (Lepidoptera:Tortricidae) on Sophora chrysophylla (Fabaceae) along an elevation gradient of dry subalpine forest on Mauna Kea, Hawaii. Pan-Pacific Entomologist 78(2): 101-109.
- British Columbia Ministry of Environment, Land and Parks. 1998a. Inventory methods for marten and weasels. Standards for Components of British Columbia's Biodiversity, No. 24, Version 2.0. Ministry of Environment, Lands and Parks, Resource Inventory Branch, Province of British Columbia, Canada Vancouver, B.C., Canada. Retrieved from http://ilmbwww.gov.bc.ca/risc/pubs/tebiodiv/marten/assets/marten.pdf.
- British Columbia Ministry of Environment, Land and Parks. 1998b. Inventory methods for small mammals: shrews, voles, mice & rats. Standards for Components of British Columbia's Biodiversity, No. 31, Version 2.0. Ministry of Environment, Lands and Parks, Resource Inventory Branch, Province of British Columbia, Canada Vancouver, B.C., Canada. Retrieved from http://ilmbwww.gov.bc.ca/risc/pubs/tebiodiv/smallmammals/assets/smlmam.pdf.
- Brown, J. M. 2008. Personal communication from Jackie Brown, Grinnell College, to Jennifer Garrison, SRGII, regarding Tephritid fly species on Mauna Kea.
- Brown, J. M., M. Todd-Thompson, A. McCord, A. O'Brien and B. O'Fallon. 2006. Phylogeny, host association, and wing pattern variation in the endemic Hawaiian fruit flies (Diptera, Tephritidae). Instrumenta Biodiversitatis VII: 1-16.
- Bruch, J. 2008. Personal communication from James Bruch, Kahoʻolawe Island Reserver Commission, to Jennfier Garrison, SRGII, regarding small mammal monitoring methods used on Kahoʻolawe.
- Bruegmann, M. M. 1996. Hawaii's dry forests. Endangered Species Bulletin 21(1): 26-27.

- Bryan, E. H., Jr. 1923. Insects from the summit of Mauna Kea. <u>Proceedings of the Hawaiian Entomological Society</u> 5(2): 287-288.
- Bryan, E. H., Jr. 1926. Additional notes on the insects occurring on Mauna Kea and Mauna Loa. Proceedings of the Hawaiian Entomological Society 6(2): 280-282.
- Bryan, L. W. 1937a. The big fence on the Big Island. Paradise of the Pacific 49(4): 15, 30.
- Bryan, L. W. 1937b. Wild sheep in Hawaii. Paradise of the Pacific 49(3): 19, 31.
- Bryan, L. W. 1939. Lake Waiau of Hawaii. Paradise of the Pacific. v. 51: p. 11.
- Bryan, L. W. 1950. Wild sheep on Mauna Kea Forest Reserve. <u>Paradise of the Pacific</u> 62(12): 112-113, 122.
- Bryan, L. W. 1973. Ahinahina. <u>Hawaiian Botanical Society Newsletter</u> 12(1): 1-2.
- Bryan, W. A. 1918. Report on the discovery of ancient glaciation on Mauna Kea, Hawaii. <u>Science</u> 47(1220): 492.
- Bryan, W. A. 1916. Mauna Loa insects. Proceedings of the Hawaiian Entomological Society 3(4): 295.
- Bryce, J. G., D. J. DePaolo, et al. 2005. Geochemical structure of the Hawaiian plume: Sr, Nd, and Os isotopes in the 2.8 km HSDP-2 section of Mauna Kea volcano. <u>Geochemistry</u>, <u>Geophysics</u>, Geosystems G super 3 **6**(9): 1-36.
- Businger, S. 2008. Personal communication from S. Businger, Professor, Department of Meteorology, University of Hawaii, Manoa to Colleen Bergmanis, SRGII.
- Businger, S., R. McLaren, et al. 2002. Starcasting. <u>Bulletin of the American Meteorological Society</u> **83**(6): 858-871.
- Businger, S., T. Cherubini and R. Lyman. 2008. Mauna Kea Weather Center (Website), University of Hawai'i Institute for Astronomy. Retrieved from http://mkwc.ifa.hawaii.edu/index.cgi.
- Byrne, D. 2008. Personal communication from D. Byrne, Manager, Mauna Kea Visitor Information Station to Kristin Duin and Colleen Bergmanis, SRGII.
- Cabin, R. J., S. G. Weller, D. H. Lorence, T. W. Flynn, A. K. Sakai, D. Sandquist and L. J. Hadway. 2000. Effects of long-term ungulate exclusion and recent alien species control preservation and restoration of a Hawaiian tropical dry forest. Conservation Biology 14(2): 439-453.
- Calanca, P. 2007. Climate change and drought occurrence in the Alpine region: How severe are becoming the extremes? Global and Plantetary Change 57(1-2): 151-160.
- Camp, R. J., T. K. Pratt, C. Bailey and D. Hu. 2006. Focal terrestrial vertebrate species: Landbirds inventory & monitoring protocol, Pacific Islands Network. Version 1.0 (draft). Inventory and Monitoring Program, Pacific Islands Network Hawaii National Park, HI.
- Cannone, N., S. Sgorbati and M. Gugfielmin. 2007. Unexpected impacts of climate change on alpine vegetation. Frontiers in Ecology and Environment 5(7): 360-364.
- Cao, G. 2007. Trade wind inversion variability, dynamics and future change in Hawai'i. <u>Geography</u> <u>Department</u>. Honolulu, HI, University of Hawaii. PhD Dissertation.
- Cao, G., T. W. Giambelluca, D. Stevens and T. Schroeder. 2007. Inversion variability in the Hawaiian trade wind regime. <u>Journal of Climate</u> 20: 1145-1160.
- Carey, E. V., M. J. Marler and R. M. Callaway. 2004. Mycorrhizae transfer carbon from a native grass to an invasive weed: evidence from stable isotopes and physiology. Plant Ecology 172: 133-141.
- Carlquist, S. 1980. Hawaii A Natural History. Lawai, Kauai, Hawaii, Pacific Tropical Botanical Garden.
- Carr, G. D. 1985. Monograph of the Hawaiian Madiinae (Asteraceae): Argyroxiphium, Dubautia, and Wilkesia. Lawai, Kauai, Hawaii, Pacific Tropical Botanical Garden.
- Carr, G. D., E. A. Powell and D. W. Kyhos. 1986. Self-incompatibility in the Hawaiian Madiinae (Compositae): An exception to Baker's Rule. Evolution 40(2): 430-434.
- Carson, H. L. and D. A. Clague 1995. Geology and biogeography of the Hawaiian Islands. Washington, DC, Smithsonian Institution Press: p. 14-28.
- Char, W. P. 1985. Botanical survey for the proposed temporary construction camp housing at Hale Pohaku, Mauna Kea, Island of Hawaii. Appendix A in: MCM Planning and University of Hawaii.

- 1985. Final supplemental environmental impact statement for construction camp housing, amendment to the Mauna Kea Science Reserve Complex Development Plan. Hale Pohaku, Hamakua, Mauna Kea, Hawaii. Prepared for MCM Planning Honolulu, HI.
- Char, W. P. 1990. Botanical Survey, JNLT and Associated Facilities for the Proposed Japanese National Large Telescope, Mauna Kea, Hawai'i. Prepared for MCM Planning Honolulu, HI.
- Char, W. P. 1999a. Botanical Resources, Hale Pohaku Mid Elevation Facilities. Appendix G of Group 70. 2000. Mauna Kea Science Reserve Master Plan. Honolulu, HI.
- Char, W. P. 1999b. Botanical Resources, Mauna Kea Summit. Appendix G of Group 70. 2000. Mauna Kea Science Reserve Master Plan. Honolulu, HI.
- Chen, Y.-L. and J. Wang. 1995b. The influences of inversion height on precipitation and airflow over the island of Hawaii. Monthly Weather Review 123(6): 1660-1676.
- Chen, Y.-L. and J.-J. Wang. 1994. Diurnal variation of surface thermodynamic fields on the island of Hawaii. Monthly Weather Review 122(9): 2125-2138.
- Chen, Y.-L. and J.-J. Wang. 1995a. The effects of precipitation on the surface temperature and airflow over the island of Hawaii. Monthly Weather Review **123**(3): 681-694.
- Chimera, C. G. 2004. Investigating seed dispersal and seed predation in a Hawaiian dry forest community: Implications for conservation and management. <u>Botany Department</u>. Honolulu, HI, University of Hawaii: 125.
- Christensen, N. L., A. M. Bartuska, et al. 1996. The Report of the Ecological Society of America Committee on the Scientific Basis for Ecosystem Management. <u>Ecological Applications</u> **6**(3): 665-691.
- Clague, D. A. 1974. The Hawaiian-Emperor seamount chain: its origin, petrology, and implications for plate tectonics. San Diego, CA, University of California: 216 p.
- Clague, D. A. 1987. Hawaiian xenolith populations, magma supply rates, and development of magma chambers. Bulletin of Volcanology **v. 49**: p. 577-587.
- Clague, D. A. 1996. The growth and subsidence of the Hawaiian-Emperor volcanic chain. A. Keast and S. E. Miller. Amsterdam, SPB Academic Publishing: p. 35-50.
- Clague, D. A. and G. B. Dalrymple 1989. Tectonics, geochronology, and origin of the Hawaiian-Emperor Chain. E. L. Winterer, D. M. Hussong and R. W. Decker. Boulder, CO, Geological Society of America. **The Geology of North America, v. N:** p. 187-217; pls. 1A, 1B, 2A, 2B, 3A, and 5 (folded maps) in separate box include Hawaii.
- Cleghorn, P. L., Dye, Tom, Weisler, Marshall, and Sinton, John. 1985. A preliminary petrographic study of Hawaiian stone adze quarries. <u>Journal of the Polynesian Society</u> **94**(3): 235-251.
- Cole, F. R., A. C. Medeiros, L. L. Loope and W. W. Zuehlke. 1992. Effects of the Argentine Ant on Arthropod Fauna of Hawaiian High-Elevation Shrubland. Ecology 73(4): 1313-1322.
- Cole, F. R., L. L. Loope, A. C. Medeiros, J. A. Raikes, C. S. Wood and L. J. Anderson. 1995b. Ecology of introduced game birds in high-elevation shrubland of Haleakala National Park. Technical Report No. 96. Cooperative National Park Resources Studies Unit, University of Hawaii at Manoa Honolulu, HI.
- Cole, F. R., L. L. Loope, A. C. Medeiros, J. A. Raikes and C. S. Wood. 1995a. Conservation implications of introduced game birds in high-elevation hawaiian shrubland. Conservation Biology 9(2): 306-313.
- Cole, I. 2007. Mauna Kea Ice Age Lake Waiau weed survey. Internal Memo. DLNR NARS.
- Collins, W. 1957. Mauna Kea hunter's paradise. Paradise of the Pacific 69(7): 16-19.
- Conant, S. 1977. The breeding biology of the Oahu 'Elepaio. Wilson Bulletin 89: 193-210.
- Conant, S. 1980. Recent records of the 'ua'u (dark-rumped petrel) and the 'a'o (Newell's shearwater) in Hawaii. 'Elepaio 41(2): 11-13.
- Conant, S., M. Hadfield, P. Aldrich, D. Barshis, N. Bustos, T. Daly-Engel, J. Eble, H. Eijzenga, J. Eijzenga, K. Hall, E. Huppman, K. Page, D. Reding, A. Stompert, N. Sugii, A. Whitehead and N. Yeung. 2004. Mauna Kea Biological Resources Management Plan. Department of Zoology, University of Hawaii Honolulu, HI.

- Cooper, A. 2008. Personal communication from Andrew Cooper, to Colleen Bergmanis, SRGII, regarding nieve penitentes on Mauna Kea.
- Copman, L. 2006. Light Pollution Solutions. Cosmic Matters (Winter 2006).
- Copman, L. 2007. The Right to Starlight: Lighting Success Stories. Cosmic Matters (Summer 2007).
- Coulter, J. W. 1939. Meteorological reports of the Mauna Kea Expedition, 1935 (III); Climate of the Humuula Plateau. <u>Bulletin of the American Meteorological Society</u> **20**: 417-419.
- Cowie, R. H., G. M. Nishida, Y. Basset and S. M. I. Gon. 1995. Patterns of land snail distribution in a montane habitat on the island of Hawaii. <u>Malacologia</u> 36(1-2): 155-169.
- Cuddihy, L. W. 1988. Vegetation zones of the Hawaiian Islands. <u>Conservation Biology in Hawaii</u>. C. P. Stone and D. B. Stone. Honolulu HI, University of Hawaii Cooperative National Park Resources Study Uni
- Cuddihy, L. W. and C. P. Stone. 1990. Alteration of native Hawaiian vegetation: effects of humans, their activities and introductions. Honolulu, Hawaii, University of Hawaii, Cooperative National Park Resources Studies Unit.
- D'antonio, C. M., J. T. Tunison, et al. 2000. Variation in the impact of exotic grasses on native plant composition in relation to fire across an elevation gradient in Hawaii. <u>Australian Journal of Ecology</u> **25**: 507-522.
- da Silva, S. C. 2006. Climatological analysis of meteorological observations at the summit of Mauna Kea. Physics Department, University of Lisbon: 77 p.
- Daehler, C. C. 2005. Upper-montane plant invasions in the Hawaiian Islands: patterns and opportunities. Perspectives in Plant Ecology, Evolution and Systematics 7: 203–216.
- Daehler, C. C. and J. Denslow. 2008. Weed risk assessment for Hawaii and Pacific Islands. from http://www.botany.hawaii.edu/faculty/daehler/wra/full_table.asp.
- Daingerfield, R. L. 1922. Lake Waiau, near summit of Mauna Kea. Paradise of the Pacific 35(12): 89.
- Daly, H. D. and K. N. Magnacca. 2003. *Insects of Hawaii: Hawaiian Hylaeus (Nesoprosopis) Bees (Hymenoptera: Apoidea)*. Honolulu, University of Hawaii Press.
- Danby, R. K. and D. S. Hik. Responses of white spruce (Picea glauca) to experimental warming at a subarctic alpine treeline. Global Change Biology 13(2): 437-451.
- Day, R. H., B. A. Cooper and R. J. Blaha. 2003. Movement patterns of Hawaiian petrels and Newell's Shearwaters on the Island of Hawaii. Pacific Science 57(2): 147-159.
- Dean, W. R. J., S. J. Milton and W. R. Siegfried. 1990. Dispersal of seeds as nest material by birds in semiarid Karoo shrubland. <u>Ecology</u> 71(4): 1299-1306.
- Debussche, M. and P. Isenmann. 1994. Bird-dispersed seed rain and seedling establishment in patchy mediterranean vegetation. Oikos 69: 414-426.
- DePaolo, D. J. and E. M. Stolper 1996. Models of Hawaiian volcano growth and plume structure: Implications of results from the Hawaii Scientific Drilling Project. Journal of Geophysical Research 101(B5): 11.643-11.654.
- Department of the Interior: Fish and Wildlife Service. 1950. Hawaii has Unique Game Management Problems, FWS Official Says.
- Dickie, E. G. 1967. Skiing on top of the world. Honolulu 1(9): 22, 23, 34-36.
- Dillon, R. D. 1979. Microbiology of an isolated alpine lake (Lake Waiau, Hawaii). <u>National Geographic Society, Research Reports</u>: 97-100.
- Division of Plant Industry. 1992. List of Plant Species Designated as Noxious Weeds for Eradication or Control Purposes by the Hawaii Department of Agriculture. Hawaii Administrative Rules, Title 4 Department Of Agriculture, Subtitle 6 Division Of Plant Industry, Chapter 68 Noxious Weed Rules.
- DLNR. 1977. The Mauna Kea Plan. <u>Mauna Kea</u>. Honolulu, HI, Hawaii Dept. of Land and Natural Resources: 12 p.
- DLNR. 1995. Revised management plan for the UH Management areas on Mauna Kea, Hamakua, Hawaii. Mauna Kea. Honolulu, HI, Hawaii Dept. of Land and Natural Resources.

- DOFAW. n.d. Hawaii's Most Invasive Horticultural Plants. Retrieved Sept. 22, 2007, from http://www.state.hi.us/dlnr/dofaw/hortweeds/specieslist.htm.
- Dorn, R. I., F. M. Phillips, et al. 1991. Glacial chronology. <u>National Geographic Research & Exploration</u> **7**(4): 456-471.
- Douglas, D. 1914. Journal, 1823-1834. London, Wm. Wesley and Son
- Duman, J. G. and S. L. Montgomery. 1991. Subzero temperature adaptations in arthropods from the summit of Mauna Kea, Hawaii. Journal of Experimental Zoology 259: 409-412.
- Ebel, B. 2000. Hydrologic modeling of Lake Waiau, Hawaii. (Student paper). St. Louis, MO, Washington University.
- Edwards, J. S. 1987. Arthropods of alpine Aeolian ecosystems. <u>Annual Review of Entomology</u> 32: 163-179.
- Edwards, J. S. 1988. Aeolian communities: Mt. Saint Helens and Mauna Kea. Wings 13(2): 8-9.
- Ehlmann, B., R. E. Arvidson, B. L. Joliff, S. S. Johnson, B. Ebel, N. Lovenduski, J. D. Morris, J. A. Byers, N. O. Snider and R. E. Criss. 2005. Hydrologic and isotopic modeling of alpine Lake Waiau, Mauna Kea, Hawaii. Pacific Science 59(1): 1-15.
- Ehses, J. 2007. Thesis Summary: GIS-gestützte Abschätzung möglicher Permafrostvorkommen am Mauna Kea, Hawaii, mittels klimageographischer Modellierung University of Aachen, Germany: 2 p.
- Eiben, J. 2008. Personal communication from Jesse Eiben, University of Hawaii, to Jennifer Garrison, SRGII, regarding current wekiu bug study.
- Eisele, J., W. Abouchami, et al. 2003. The 320 kyr Pb isotope evolution of Mauna Kea lavas recorded in the HSDP-2 drill core. G³ (Geochemistry, Geophysics, Geosystems) v. 4(no. 5; 8710): doi:10.1029/2002GC000339.
- Ekern, P. C. and J. H. Chang. 1985. Pan evaporation: State of Hawaii, 1894-1983, Department of Land and Natural Resources, State of Hawaii. **R74:** 172.
- Eldridge, D., S. Skinner and T. J. Entwisle. 2003. Survey guidelines for non-vascular plants. Botanic Gardens Trust Sydney, Australia. Retrieved from http://www.environment.nsw.gov.au/resources/nature/SurveyGuideNon-VascularPlants.pdf.
- Elzinga, C. L., D. W. Salzer and J. W. Willoughby. 1998. Measuring and monitoring plant populations. Technical Reference 1730–1. Bureau of Land Management Denver, CO.
- Englund, R.A., D.J. Preston, A.E. Vorsino, S. Meyers, and L. L. Englund. 2009. Results of the 2007–2008 Invasive Species and Wekiu Bug (Nysius wekiuicola) Surveys on the Summit of Mauna Kea, Hawai'i Island. April 2009 Draft Report. Hawaii Biological Survey Report prepared for the Office of Mauna Kea Management. 73 pp.
- Englund, R. A. and D. J. Preston. 2008. Personal communications from Ron Englund and David Preston, Bishop Museum, to Jennifer Garrison, SRGII, regarding invertebrate species found at Lake Waiau.
- Englund, R. A., D. A. Polhemus, F. G. Howarth and S. L. Montgomery. 2002. Range, habitat, and ecology of the wekiu bug (Nysius wekiuicola), a rare insect species unique to Mauna kea, Hawai`i Island. Final report. Prepared for Office of Mauna Kea Management, University of Hawaii. Hawaii Biological Survey Report 2002-023. Bishop Museum Honolulu.
- Englund, R. A., A. E. Vorsino and H. M. Laederich. 2007. Results of the 2006 Wekiu Bug (Nysius wekiuicola) Surveys on Mauna Kea, Hawai'i Island. Final Report. Prepared for Office of Mauna Kea Management. Hawaii Biological Survey Report 2007-003. Hawaii Biological Survey, Bishop Museum Honolulu, HI.
- Englund, R. A., A. E. Vorsino, H. M. Laederich, A. Ramsdale and M. McShane. 2006. Results of 2005 Wekiu Bug (Nysius wekiuicola) Surveys on Mauna Kea, Hawai'i Island. Final Report. Prepared for Office of Mauna Kea Management. Hawaii Biological Survey Report 2006-010. Hawaii Biological Survey, Bishop Museum Honolulu, HI.
- Englund, R. A., A. Ramsdale, M. McShane, D. J. Preston, S. Miller and S. L. Montgomery. 2005. Results of 2004 Wekiu Bug (Nysius wekiuicola) Surveys on Mauna Kea, Hawai'i Island. Final Report.

- Prepared for Office of Mauna Kea Management. Hawaii Biological Survey Report 2005-003. Hawaii Biological Survey, Bishop Museum Honolulu, HI.
- Erschbamer, B. 2007. Winners and losers of climate change in a central alpine glacier foreland. <u>Arctic Antarctic and Alpine Research</u> 39(2): 237-244.
- Evans, S. 2008. Personal communication from Steve Evans, PTA, to Jennifer Garrison, SRGII, regarding invasive species management activities at PTA.
- Exley, R. A., D. P. Mattey, et al. 1986. Carbon isotope systematics of a mantle "hotspot": a comparison of Loihi Seamount and MORB glasses. <u>Earth and Planetary Science Letters</u> v. 78: p. 189-199.
- Fan, P.-F. 1978. Mineralogy of sediments from Lake Waiau, Hawaii. Pacific Science 32(2): 219-221.
- Fancy, S. G. and C. J. Ralph. 1997. 'Apapane (Himatione sanguinea). <u>The Birds of North America, No. 296</u>. A. Poole and F. Gill, The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, D.C.: 16 p.
- Fancy, S. G. and C. J. Ralph. 1998. 'I'iwi (Vestiaria coccinea). <u>The Birds of North America, No. 327</u>. A. Poole and F. Gill. Philadelphia, PA, The Birds of North America, Inc.
- Feigenson, M. D., L. L. Bolge, et al. 2003. REE inverse modeling of HSDP2 basalts: evidence for multiple sources in the Hawaiian plume. G³ (Geochemistry, Geophysics, Geosystems) v. 4(no. 2, 8706, doi:10.1029/2001GC000271): 25 p.
- Finnamore, A. T., N. N. Winchester and V. M. Behan-Pelletier. 2002. Protocols for measuring biodiversity: Arthropod monitoring in terrestrial ecosystems. Environment Canada Ecological Monitoring and Assessment Network (EMAN). Retrieved from http://www.eman-rese.ca/eman/ecotools/protocols/terrestrial/arthropods/intro.html.
- Flenley, J. R. 1998. Tropical forests under the climates of the last 30,000 years. <u>Climatic Change</u> 39: 177–197.
- Fodor, R. V. 2001. The role of tonalite and diorite in Mauna Kea Volcano, Hawaii, magmatism: petrology of summit-region leucocratic xenoliths. <u>Journal of Petrology</u> **42**(9): 1685-1704.
- Forsyth, S. A. 2002a. Demographic modeling of Hawaiian silverswords, and its implications for conservation, University of Arizona: 283 p.
- Fox, S. and IfA. 2007. A guide to the plants of Mauna Kea. <u>Mauna Kea Visitor Information Station (web page)</u>.
- Frasher, H. R., V. Parker-Geisman and G. R. Parish. 2007. Hawaiian hoary bat inventory in National Parks on the islands of Hawai'i, Maui, and Moloka'i. Technical Report no.140. Pacific Cooperative Study Unit, Dept of Botany, University of Hawaii Honolulu, HI.
- Frear, W. F. 1913. Waiau lake is the only lake in Hawaii on which ice skating may be enjoyed. Mid-Pacific Magazine 6(5): 482.
- Freed, L. A., Cann, Rebecca L., Goff, M. Lee, Kuntz, Wendy A., and Bodner, Gustav R. 2005. Increase in avian malaria at upper elevation in Hawaii. <u>Condor</u> 107(4): 753-764.
- Freed, L. A., S. Conant and R. C. Fleischer. 1987. Evolutionary ecology and radiation of Hawaiian passerine birds. <u>Trends in Ecology and Evolution</u> 2: 196-203.
- Fretz, J. S. 2008. Personal communication from Scott Fretz, DLNR, to Jennifer Garrison, SRGII, regarding management of DLNR lands on Mauna Kea.
- Frey, F. A., Wise, W.S., Garcia, M.O., West, H., Kwon, S.T., and Kennedy, A. 1990. Evolution of Mauna Kea volcano, Hawaii: Petrologic and geochemical constraints on postshield volcanism. <u>Journal of Geophysical Research</u> **95**(B2): 1271-1300.
- Friar, E. A., T. Ladoux, E. H. Roalson and R. H. Robichaux. 2000. Microsatellite analysis of a population crash and bottleneck in the Mauna Kea silversword, Argyroxiphium sandwicense ssp. Sandwicense (Asteraceae), and its implications for reintroduction. Molecular Ecology 9(12): 2027-2034.
- Furumoto, A. S., W. A. Wiebenga, et al. 1973. Crustal structure of the Hawaiian Archipelago, northern Melanesia, and the central Pacific Basin by seismic refraction methods. <u>Tectonophysics</u>. S. Mueller. v. 20: p. 153-164.

- Gagne, W. C. 1986. Nysius wekiuicola. Proceedings of the Hawaiian Entomological Society 26: 7.
- Gagne, W. C. and C. C. Christensen. 1985. Conservation status of native terrestrial invertebrates in Hawai'i. <u>Hawaii's terrestrial ecosystems: preservation and management. Proceedings of a symposium held June 5-6, 1984 at Hawaii Volcanoes National Park</u>. C. P. Stone and M. J. Scott. Honolulu, HI, Cooperative National Park Resources Studies Unit, University of Hawaii: 105-125.
- Gagne, W. C. and F. G. Howarth. 1982. Entomologists discover new life. <u>Ka Elele Bishop Museum</u> Newsletter 9(2): 1,6.
- Gagné, W. C. and L. W. Cuddihy. 1990. Vegetation. Manual of the Flowering Plants of Hawai'i. W. L. Wagner, D. R. Herbst and S. H. Sohmer. Honolulu, Bishop Museum Press: 45-114.
- Gambino, P., A. C. Medeiros and L. L. Loope. 1987. Introduced vespids Paravespula pensylvanica prey on Maui's endemic arthropod fauna. <u>Journal of Tropical Ecology</u> 3: 169-170.
- Gambino, P., A. C. Medeiros and L. L. Loope. 1990. Invasion and colonization of upper elevations on East Maui (Hawaii) by Vespula pensylvanica (Hymenoptera: Vespidae). <u>Annals of the Entomological Society of America 83(6)</u>: 1088-1095.
- Gardner, D. E. 1997. Botryosphaeria mamane sp. nov. associated with witches'-brooms on the endemic forest tree Sophora chrysophylla in Hawaii. Mycologia 89(2): 298-303.
- Garrison, J. S. E. 2003. The role of alien tree plantations and avian seed-dispersers in native dry forest restoration in Hawai'i. Zoology Department. Honolulu, HI, University of Hawaii: 370 p.
- Gemma, J. N. and R. E. Koske. 2001. Use of mycorrhizae in restoration of Hawaiian habitats. Retrieved 9/11/02, 2002, from http://www.hawaii.edu/scb/scinativ_mycor.html.
- Gemma, J. N., R. E. Koske and M. Habte. 2002. Mycorrhizal dependency of some endemic and endangered Hawaiian plant species. <u>American Journal of Botany</u> 89(2): 337–345.
- Gerrish, G. 1979. Vegetation Survey. Appendix B in: Group 70. 1980. Revised environmental impact statement for Hale Pohaku: mid-elevation facilities master plan, Hamakua, Mauna Kea, Hawaii. Prepared for Group 70 Honolulu, HI.
- Giambelluca, T. W. 2008. Personal communication from Tom Giambelluca, University of Hawaii Geography Department, to Jennifer Garrion, SRGII, regarding climate change predictions in Hawai'i. Honolulu, HI.
- Giambelluca, T. W. and M. S. A. Luke. 2007. Climate change in Hawai'i's mountains. <u>Cirmount (The Newsletter of the Consortium for Integrated Climate Research in Western Mountains)</u> 1(2): 13-18.
- Giambelluca, T. W. and M. Sanderson. 1993. The Water Balance and Climate Classification. <u>Prevailing Trade Winds, Weather and Climate in Hawai'i.</u> M. Sanderson. Honolulu, University of Hawaii Press.
- Giffin, J. G. 1982. Ecology of the mouflon sheep on Mauna Kea: final report. Pittman-Robertson Proj. W-17-R, Study No. R-III, 1975-1979. State of Hawaii, Dept. of Land and Natural Resources, Division of Forestry and Wildlife.
- Gilbertson, R. L., D. E. Desjardin, J. D. Rogers and D. E. Hemmes. 2001. Fungi from the Mamane-Naio vegetation zone of Hawaii. <u>Fungal Diversity</u>; <u>An International Journal of Mycology</u> 6: 35-68.
- Gillies, C. and D. Williams. 2003. Using tracking tunnels to monitor rodents and mustelids. Threats Science Team, Terrestrial Conservation Unit, Department of Conservation Hamilton, New Zealand.
- Gon, S. 2003. Applications of Traditional Ecological Knowledge and Practices of Indigenous Hawaiians to the Revegetation of Kahoolawe. Ethnobotany Research & Applications 1: 5-20.
- Good, L. 2003. Mauna Kea's Visitor Information Station. Na Kilo Hoku: The Ones who Look to the Stars Volume, DOI:
- Goodrich, J. 1826. Notice of the volcanic character of the Islands of Hawaii. <u>American Journal of Science</u> 11(Series 1): 2-7.
- Goodrich, J. 1833a. Climbing Mauna Kea in "Letters to the Editor". <u>American Journal of Science</u> 25: 199-201.

- Goodrich, J. 1833b. Notices of some of the volcanos and volcanic phenomena of Hawaii, (Owyhee,) and other islands in that group, in a letter from Mr. Joseph Goodrich, missionary, dated Nov. 17, 1832. Amercican Journal of Science 25: 199-203.
- Goodrich, J. 1826. Letters about volcanoes of Hawaii. American Journal of Science 11: 2-36.
- Gorresen, M., A. Miles, C. Todd, F. Bonaccorso and T. Weller. 2008. Assessing bat detectability and occupancy with multiple automated echolocation detectors. <u>Journal of Mammalogy</u> 89(1): 11-17.
- Gray, E. M., P. C. Banko, S. J. Dougill, D. Goltz, L. Johnson, M. E. Laut, J. D. Semones and M. R. Wiley. 1999. 1998 Palila population census; breeding and nonbreeding censuses of the 1998 palila population on Mauna Kea, Hawaii. <u>Elepaio</u> 59(5): 33-39.
- Gregory, H. E. and C. K. Wentworth 1937. General features and glacial geology of Mauna Kea, Hawaii. Geological Society of America Bulletin 48: 1719-1942.
- Griffin, C. R., P. W. C. Paton and T. S. Baskett. 1998. Breeding ecology and behavior of the Hawaiian hawk. <u>The Condor</u> 100(4): 654-662.
- Group 70. 1980. Hale Pohaku, Hamakua, Mauna Kea, Hawaii, mid-elevation facilities master plan: complex development report. <u>Infrastructure</u>. Honolulu, HI, Dept. of Land and Natural Resources, State of Hawaii.
- Group 70. 1982. Mauna Kea Science Reserve complex development plan: draft environmental impact statement, Mauna Kea, Hamakua, Hawaii. Honolulu, HI, Prepared for Research Corporation of the University of Hawaii. Vol. I, Vol. II.
- Group 70. 1983a. Mauna Kea Science Reserve complex development plan, Mauna Kea, Hamakua, Hawaii. Mauna Kea. Honolulu, HI, Prepared for Research Corporation of the University of Hawaii.
- Group 70. 1983b. Mauna Kea Science Reserve complex development plan: final environmental impact statement, Mauna Kea, Hamakua, Hawaii. Mauna Kea. Honolulu, HI, Prepared for Research Corporation of the University of Hawaii.
- Group 70. 1987. Mauna Kea Science Reserve complex development plan, Mauna Kea, Hamakua, Hawaii. 1983. Amended May 1987. <u>Mauna Kea</u>. Honolulu, HI, Prepared for Research Corporation of the University of Hawaii.
- Group 70 International. 1999. Mauna Kea Science Reserve Master Plan: final environmental impact statement. Master Plan. Honolulu, HI, Prepared for University of Hawaii. Vol.I, Vol.II.
- Group 70 International 2000. Mauna Kea Science Reserve Master Plan. Adopted by the University of Hawaii Board of Regents on June 16, 2000. <u>Master Plan</u>. Honolulu, HI, Prepared for the University of Hawaii. **Vol.I, Vol.II**.
- Grumbine, R. E. 1994. What is ecosystem management? Conservation Biology 8(1): 27-38.
- Gruner, D. S. and D. Foote. 2000. Management Strategies for Western Yellowjackets in Hawaii. 6 p.
- Guinness, E. A., R. E. Arvidson, B. L. Jolliff, K. S. Deal, F. P. Seelos, R. V. Morris, D. W. Ming and T. G. Graff. 2003. Mapping hydrothermal alteration zones on Mauna Kea using AVIRIS data: an analog for Mars [abs.]. <u>Eos, Transactions, American Geophysical Union supp.</u> v. 84(no. 46): p. F934-F935.
- Guppy, H. B. 1897. The summit of Mauna Loa. Nature 57(1462): 20-21.
- Habte, M. 2000. Mycorrhizal fungi and plant nutrition. <u>Plant Nutrient Management in Hawaii's Soils</u>, <u>Approaches for Tropical and Subtropical Agriculture</u>. J. A. Silva and R. Uchida. Honolulu, HI, College of Tropical Agriculture and Human Resources, University of Hawaii at Manoa: 127-131.
- Hadway, L. 2007. Personal communication from Lisa Hadway, DLNR, to Kristin Duin, SRGII, regarding conditions at Ice Age NAR.
- Hadway, L. and M. G. Hadfield. 1998. Final Report: Assessment of the current status of Partulina tree-snail species on the Island of Hawai'i with comments on additional land snail species. Department of Zoology, University of Hawaii. Prepared for the U.S. Fish and Wildlife Service Honolulu, HI.
- Hadway, L. and M. Hadfield. 1999. Conservation status of tree snail species in the genus Partulina (Achatinellinae) on the Island of Hawai'i: A modern and historical perspective. <u>Pacific Science</u> 53(1): 1-14.

- Hadway, L. J. 2008. Personal communication from Lisa Hadway, DLNR, to Jennifer Garrison, SRGII, regarding conditions at Ice Age NAR and MKSR.
- Hamilton, K. 2007a. Interim report on results from PRIDE Project "Global Warming Effects on Extreme Weather Events in Hawaii Studied Using Results from Ultrafine Resolution Global Models" Honolulu, HI, International Pacific Research Center (IPRC), University of Hawaii: 10 p.
- Hamilton, K. 2007. Personal communication from Kevin Hamilton, International Pacific Research Center (IPRC), University of Hawaii, to Jennifer Garrison, SRGII, regarding climate change models.
- Hansen, D. M., J. M. Olesen and C. G. Jones. 2002. Trees, birds and bees in Mauritius: exploitative competition between introduced honey bees and endemic nectarivorous birds? <u>Journal of Biogeography</u> 29: 721–734.
- Hartt, C. E. and M. C. Neal. 1940. The plant ecology of Mauna Kea, Hawaii. Ecology 21(2): 237-266.
- Hawaii Ant Group. 2007. A plan for prevention of establishment of new ant species in Hawaii, with special attention to the red imported fire ant (*Solenopsis invicta*) and little fire ant (*Wasmannia auropunctata*): 51 p.
- Hawaii Audubon Society. 1997. Hawaii's Birds. Honolulu, HI, Hawaii Audubon Society.
- Hawaii Biodiversity and Mapping Program. n.d. Cystopteris douglasii, Douglas' Bladder Fern. Retrieved November 12, 2007, from http://hbmp.hawaii.edu/printpage.asp?spp=PPDRY07020.
- Hawaii State Office of Environmental Quality Control. 2008. Madagascar fireweed biocontrol project (*Secusio extensa*) (HRS 343-5(b) DEA). The Environmental Notice (April 23, 2008): 11.
- Hawkes, C. V., J. Belnap, C. D'Antonio and M. K. Firestone. 2006. Arbuscular mycorrhizal assemblages in native plant roots change in the presence of invasive exotic grasses. <u>Plant and Soil</u> 281: 369-380.
- Hayashi, S. 2008. Personal communication from Saeko Hayashi, Subaru Telescope, to Andrew Hood, SRGII, regarding collection of climate data.
- HaySmith, L. 2008. Personal communication from Leslie HaySmith, National Park Service, to Jennifer Garrison, SRGII, regarding status of Pacific Islands Network (NPS) monitoring protocols.
- HaySmith, L., F. L. Klasner, S. H. Stephens and G. H. Dicus. 2005. Pacific Island Network vital signs monitoring plan. Natural Resource Report NPS/PACN/NRR—2006/003. National Park Service Fort Collins, CO. Retrieved from http://science.nature.nps.gov/im/monitor/plans/PACN MonitoringPlan.pdf.
- Hemmes, D. E. and D. E. Desjardin. 2002. Mushrooms of Hawai'i. Berkeley, CA, Ten Speed Press.
- Hess, S. C., P. C. Banko, et al. 1999. Factors related to the recovery of subalpine woodland on Mauna Kea, Hawaii. <u>Biotropica</u> **31**(2): 212-219.
- Hess, S. C., P. C. Banko, G. J. Brenner and J. D. Jacobi. 1996. Tree densities in a subalpine woodland on Mauna Kea, Hawaii. <u>Proceedings of the Western Association of Fish and Wildlife Agencies: Honolulu</u> Hawaii, July 22 26, 1996.: 225-234.
- Hess, S. C., P. C. Banko, G. J. Brenner and J. D. Jacobi. 1999. Factors related to the recovery of subalpine woodland on Mauna Kea, Hawaii. <u>Biotropica</u> 31(2): 212-219.
- Hess, S. C., P. C. Banko, M. H. Reynolds, G. J. Brenner, L. P. Laniawe and J. D. Jacobi. 2001. Drepanidine movements in relation to food availability in subalpine woodland on Mauna Kea, Hawaii. Studies in Avian Biology 22: 154-163.
- Higashino, J. 2008. Personal communication from Jennifer Higashino, DOFAW, DLNR, to Jennifer Garrison, SRGII, regarding presence of invasive fountain grass on Mauna Kea and Mauna Loa.
- Hitchcock, A. S. 1917a. A botanical trip to the Hawaiian Islands. The Scientific Monthly 5(4): 323-349.
- Hitchcock, A. S. 1917b. A botanical trip to the Hawaiian Islands. II. <u>The Scientific Monthly</u> **5**(5): 419-432.
- Hoffman, R. L., T. J. Tyler, G. L. Larson, M. J. Adams, W. Wente and S. Galvan. 2005. Sampling protocol for monitoring abiotic and biotic characteristics of mountain ponds and lakes. Retrieved from http://science.nature.nps.gov/im/monitor/refer.cfm?Protocol_ID=170.

- Hofmann, A. W. and K. P. Jochum. 1996. Source characteristics derived from very incompatible trace elements in Mauna Loa and Mauna Kea basalts, Hawaii Scientific Drilling Project. <u>Journal of Geophysical Research</u>. v. 101: p. 11,831-11,839.
- Holcomb, R. T., B. K. Nelson, et al. 2000. Overlapping volcanoes: the origin of Hilo Ridge, Hawaii. Geology v. 28(no. 6): p. 547-550.
- Holt, A. 1996. An alliance of biodiversity, agriculture, health, and business interests for improved alien species management in Hawaii. Presented at the Norway/UN Conference on Alien Species, July 1-5, 1996. Honolulu, HI, The Nature Conservancy.
- Ho'akea LLC dba Ku'iwalu. 2009. Mauna Kea Comprehensive Management Plan: UH Management Areas.
- Hotchkiss, S. and J. O. Juvik. 1999. A Late-Quaternary Pollen Record from Ka'au Crater, O'ahu, Hawai'i. Quaternary Research 52: 115–128.
- Hotchkiss, S., P. M. Vitousek, et al. 2000. Climate cycles, geomorphological change, and the interpretation of soil and ecosystem development. <u>Ecosystems</u> **3**(6): 522-533.
- Howarth, F. G. 1983. A brachypterous Nysius sp. from the summit of Mauna Kea. <u>Proceedings of the Hawaiian Entomological Society</u> 24(2 & 3): 160-161.
- Howarth, F. G. 1985. Impacts of alien land arthropods and mollusks on native plants and animals in Hawai'i. <u>Hawaii's terrestrial ecosystems: preservation and management</u>. C. P. Stone and J. M. Scott. Honolulu, University of Hawaii Press: 149-179.
- Howarth, F. G. 1987. Evolutionary ecology of aeolian and subterranean habitats in Hawaii. <u>Trends in Ecology and Evolution</u> 2: 220-223.
- Howarth, F. G. and F. D. Stone. 1982. An assessment of the arthropod fauna and aeolian ecosystem near the summit of Mauna Kea, Hawaii. Submitted to Group 70. Bernice P. Bishop Museum Honolulu.
- Howarth, F. G. and S. L. Montgomery. 1980. Notes on the ecology of the high altitude aeolian zone on Mauna Kea. <u>Elepaio</u> 41(3): 21-22.
- Howarth, F. G., G. J. Brenner and D. J. Preston. 1999. An arthropod assessment within selected areas of the Mauna Kea Science Reserve. Final Report. Prepared for the University of Hawaii Institute of Astronomy. Appendix J in Group 70. 2000. Mauna Kea Science Reserve Master Plan. Bishop Museum and Pacific Analytics Honolulu.
- Hu, D., C. Glidden, J. S. Lippert, L. Schnell, J. S. MacIvor and J. Meisier. 2001. Habitat use and limiting factors in a population of Hawaiian dark-rumped petrels on Mauna Loa, Hawaii. <u>Studies in Avian</u> Biology 22: 234-242.
- Hughes, F., P. M. Vitousek and T. Tunison. 1991. Alien grass invasion and fire in the seasonal submontane zone of Hawaii. Ecology 72(2): 743-747.
- Hughes, M. and A. Morrison-Saunders 2002. Impact of Trail-side Interpretive Signs on Visitor Knowledge. Journal of Ecotourism 1(2&3): 122-132.
- Hutchinson, J., D. Maynard and L. Geiser. 1996. Air quality and lichens A literature review emphasizing the Pacific Northwest, USA. USDA Forest Service, Pacific Northwest Region Air Resource Management Program Portland, OR. Retrieved from http://www.fs.fed.us/r6/aq/lichen/almanac.htm#Physiological%20Responses.
- Jacobi, J. D., R. K. Loh, A. Ainsworth and P. Berkowitz. 2007. Focal terrestrial plant communities inventory & monitoring protocol, Pacific Island Network. Draft Version 1.00 (October 2007). Inventory and Monitoring Program, Pacific Islands Netweork Hawaii National Park, HI.
- Jaggar, T. A. 1925. Geological notes on Mauna Kea. <u>Hawaiian Volcano Observatory Monthly Bulletin</u> 13(10): 75-77.
- James, H. F. 2004. The osteology and phylogeny of the Hawaiian finch radiation (Fringillidae: Drepanidini), including extinct taxa. Zoological Journal of the Linnean Society 141: 207-255.
- Jasper, C. 2008. Personal communication from C. Jasper, Natural Resource Conservation Service Soil Scientist, Honolulu Office to Colleen Bergmanis, SRGII.

- Johnson, E. 2008. Personal communication from E. Johnson, Wildlife Biologist and State Hunting Coordinator, Hawaii Division of Forestry and Wildlife to Colleen Bergmanis SRGII.
- Johnson, O. W., A. J. Bennett, L. Alsworth, III., L. A. Bennett, P. M. Johnson, J. R. Morgart and R. J. Kienholz. 2001. Radio-tagged Pacific golden-plovers: the Hawaii-Alaska link, spring destinations, and breeding season survival. <u>Journal of Field Ornithology</u> 72(4): 537-546.
- Johnson, S. 2001. Modeling Lake Waiau: a study of stable isotope distribution and water chemistry (student paper). St. Louis, MO, Washington University.
- Judd, C. S. 1936. Hawaii's forests winning their battle with wild animals. <u>Honolulu Star Bulletin</u>. Honolulu, HI: March 21, Section 3: 1,7.
- Juvik, J. O. and S. P. Juvik. 1984. Mauna Kea and the myth of multiple use endangered species and mountain management in Hawaii. <u>Mountain Research and Development</u> 4(3): 191-202.
- Juvik, J. O. and S. P. Juvik. 1992. Mullein (Verbascum thapsus): The spread and adaptation of a temperate weed in the montane tropics. <u>Alien Plant Invasions in Native Ecosystems of Hawai'i:</u> <u>Management and Research C. P. Stone, C. W. Smith and J. T. Tunison. Honolulu, University of Hawaii Cooperative National Park Resources Studies Unit pp 254-270.</u>
- Juvik, S. P. and J. O. Juvik, Eds. 1998. Atlas of Hawaii. Honolulu, University of Hawaii Press.
- Kanda, H. and M. Inoue. 1994. Ecological monitoring of moss and lichen vegetation in the Syowa Station Area, Antarctica. <u>Proc. NIPR Symp. Polar Biol.</u> 7: 221-231.
- Kauahikaua, J., T. Hildenbrand, et al. 2000. Deep magmatic structures of Hawaiian volcanoes, imaged by three-dimensional gravity models. <u>Geology</u> **v. 28**(no. 10): p. 883-886.
- Kawashima, I. 2008. Personal communication from I. Kawashima, Hawaii Division of Forestry and Wildlife to Colleen Bergmanis SRGII.
- Kazakis, G., D. Ghosn, I. Vogiatzakis and V. Papanastasis. 2007. Vascular plant diversity and climate change in the alpine zone of the Lefka Ori, Crete. Biodiversity and Conservation 16(6): 1603-1615.
- Kern, M. D. and C. van Riper, III. 1984. Altitudinal variations in nests of the Hawaiian honeycreeper Hemignathus virens virens. <u>Condor</u> 86(4): 443-454.
- Kessler, C. C. 2002. Eradication of feral goats and pigs and consequences for other biota on Sarigan Island, Commonwealth of the Northern Mariana Islands. <u>Turning the tide: the eradication of invasive species</u>. C. R. Veitch and M. N. Clout. Gland, Switzerland and Cambridge, UK, IUCN SSC Invasive Species Specialist Group: 132-140.
- Kihara, S. 2008. Personal communication from S. Kihara, Environmental Health Specialist, State Department of Health Clean Air Branch regarding air quality monitoring at Mauna Kea. C. B. SRGII.
- Kjargaard, M. S. 1988. Faunal survey for the proposed VLBA antenna facility, Mauna Kea, Hamakua, Hawaii. Appendix F in MCM Planning. 1988. Final supplemental environmental impact statement VLBA Antenna Facility, Mauna Kea, Hamakua, Hawaii, September, 1988. Amendment to the Mauna Kea Science Reserve Complex Development Plan. Prepared for The National Radio Astronomy Observatory, Socorro, New Mexico.
- Klavitter, J. L., J. M. Marzluff and M. S. Vekasy. 2003. Abundance and demography of the Hawaiian hawk: Is delisting warrarnted? <u>Journal of Wildlife Management</u> 67(1): 165-176.
- Klein, F. W. 1995. Seismic hazard on the Island of Hawaii from probabilistic peak ground acceleration [abs.]. Eos, Transactions, American Geophysical Union supp. v. 76(no. 46): p. 360.
- Klein, F. W. and T. L. Wright. 2000. Catalog of Hawaiian earthquakes, 1823-1959. <u>U.S. Geological Survey Professional Paper 1623</u>: 90 p.
- Klein, F. W., A. D. Frankel, et al. 2000. Seismic-hazard maps for Hawaii. <u>U.S. Geological Survey</u> Geologic Investigations Series Map I-2724.
- Klein, F. W., A. D. Frankel, et al. 2001. Seismic hazard in Hawaii: high rate of large earthquakes and probabilistic ground-motion maps. <u>Bulletin of the Seismological Society of America</u> **v. 91**(no. 3): p. 479-498.

- Koehler, R. 2008. Personal communication from R. Koehler, Director, Mauna Kea Observatory Support Services to Colleen Bergmanis, SRGII.
- Koehler, R. 2008. Personal communication from R. Koehler, Director, Mauna Kea Observatory Support Services to Colleen Bergmanis, SRGII.
- Koske, R. E. and J. N. Gemma. 2002. Mycorrhizal status of two Hawaiian plant species (Asteraceae) in a tropical alpine habitat: The threatened Haleakala silversword (*Argyroxiphium sandwicense* subsp. *macrocephalum*) and the endemic *Dubautia menziesii*. Pacific Science 56(4): 423-430.
- Koske, R. E., J. N. Gemma and L. Englander. 1990. Vesicular-Arbuscular Mycorrhizae in Hawaiian Ericales. <u>American Journal of Botany</u> 77(1): 64-68.
- Koske, R. E., J. N. Gemma and T. Flynn. 1992. Mycorrhizae in Hawaiian Angiosperms: A survey with implications for the origin of the native flora. <u>American Journal of Botany</u> 79(8): 853-862.
- Kramer, R. J. 1971. Selected chapters: The feral cattle, Bos taurus, p.271-282; The feral goat, Capra hircus, p. 283-298; The mouflon, Ovis musimon, p. 299-313; The feral sheep, Ovis aries, p. 314-327. <a href="https://doi.org/10.1001/jaure.1001/jaure.1001/ja
- Krushelnycky, P. D., S. M. Joe, A. C. Medeiros, C. C. Daehler and L. L. Loope. 2005. The role of abiotic conditions in shaping the long-term patterns of a high-elevation Argentine ant invasion. <u>Diversity and Distributions</u> 11(4): 319-331.
- Kullman, L. 2006. Long-term geobotanical observations of climate change impacts in the Scandes of West-Central Sweden. <u>Nordic Journal of Botany</u> 24(4): 445-467.
- Landgraf, L. K. 1973. Mauna Kea and Mauna Loa silversword: alive and perpetuating. <u>Pacific Tropical Botanical Garden Bulletin</u> **3**(4): 64-66.
- Lane, R. B. 1956. Maunakea's open pit basalt mines: evaluation of archaeological Hochi-sponsored field trip. <u>Hawaii Hochi</u> **31**(176): 1.
- Lang, L. and D. Byrne. 2005. <u>Mauna Kea: A Guide to Hawai'i's Sacred Mountain</u>. Honolulu, Watermark Publishing.
- Laws, E. A. and A. H. Woodcock 1981. Hypereutrophication of an Hawaiian alpine lake. <u>Pacific Science</u> **35**(3): 257-261.
- Le Roux, J. J., A. M. Wieczorek, M. M. Ramadan and C. T. Tran. 2006. Resolving the native provenance of invasive fireweed (Senecio madagascariensis Poir.) in the Hawaiian Islands as inferred from phylogenetic analysis. Diversity and Distributions 12(6): 694–702.
- Leary, J. K., N. V. Hue, P. W. Singleton and D. Borthakur. 2005. The major features of an infestation by the invasive weed legume gorse (Ulex europaeus) on volcanic soils in Hawaii. <u>Biology and Fertility of Soils</u> 42(3): 215-223.
- Levey, D. J. 1986. Methods of seed processing by birds and seed deposition patterns. <u>Frugivores and seed dispersal</u>. A. Estrada and T. H. Fleming. Dordrecht, Dr W Junk Publishers: 147-158.
- Lewin, V. and G. Lewin. 1984. The Kalij pheasant, a newly established game bird on the island of Hawaii. Wilson Bulletin 96: 634-646.
- Lewis, D. C. 1937. Skiing under the Hawaiian skies. Paradise of the Pacific 49(6): 17, 31.
- Lippiatt, S. 2005. The isolation and identification of diatoms from Lake Waiau sediments. <u>Journal of</u> Young Investigators **13**(4): 6 p.
- Lockwood, J. 2000. Mauna Kea Science Reserve Geological Resources Management Plan. Appendix H of Group 70. 2000. Mauna Kea Science Reserve Master Plan. Honolulu, HI.
- Loh, R. K., A. Ainsworth, B. Miner, R. Russell, J. Makaike and J. T. Tunison. 2000. Mullein survey and removal efforts on Mauna Loa in Hawai'i Volcanoes National Park. Technical Report no.126. Pacific Cooperative Study Unit, Dept of Botany, University of Hawaii Honolulu, HI.
- Loope, L. L. and T. W. Giambelluca. 1998. Vulnerability of island tropical montane cloud forests to climate change, with special reference to East Maui, Hawaii. <u>Climatic Change</u> 39: 503–517.
- Loope, L. L., F. G. Howarth, F. Kraus and T. K. Pratt. 2001. Newly emergent and future threats of alien species to Pacific birds and ecosystems. <u>Studies in Avian Biology</u> 22: 291-304.

- Lyons, C. J. 1875. Land matters in Hawaii. No. 2. The Islander 1(19): 111.
- MacDonald, G. A. 1945. Ring structures at Mauna Kea, Hawaii. <u>American Journal of Science</u> **243**(4): 210-217.
- Macdonald, G. A. and A. T. Abbott. 1970. Volcanoes in the sea. Honolulu, University of Hawaii Press.
- Macdonald, G. A., A. T. Abbott, et al. 1983. <u>Volcanoes in the sea: the geology of Hawaii</u>. Honolulu, University of Hawaii Press.
- Mace, G. M. and A. Purvis. 2007. Evolutionary biology and practical conservation: bridging a widening gap. <u>Molecular Ecology</u>.
- Maciolek, J. A. 1969. Fresh water lakes in Hawaii. <u>Verhandlungen-Internationale Vereinigung fur</u> theoretische und angewandte Limnologie (Proceedings-International Association of Theoretical and <u>Applied Limnology</u>) **17**: 386-391.
- Magnacca, K. N. 2007. Conservation status of the endemic bees of Hawai'i, Hylaeus (Nesoprosopis) (Hymenoptera: Colletidae). <u>Pacific Science</u> 61(2): 173–190.
- Magnacca, K. N. 2008. Personal communication from Karl Magnacca to Jennifer Garrison, SRGII, regarding native bees at Hale Pohaku and in the Mauna Kea Science reserve.
- Maly, K. 1999. 1999 Mauna Kea Science Reserve and Hale Pohaku Complex Development Plan Update: Oral History and Consultation Study and Archival Literature Research. Ahupua'a of Ka'aho (Hamakua District) and Humu'ula (Hilo District), Island of Hawai'i (various TMK). Appendix I of Group 70. 2000. Mauna Kea Science Reserve Master Plan. Honolulu, HI.
- Maly, K. and O. Maly 2005. Mauna Kea, ka piko kaulana o kaaina: Mauna Kea, the famous summit of the land. Hilo, HI, Kumu Pono Associates, LLC: 650 p.
- Markin, G. P., L. A. Dekker, J. A. Lapp and R. F. Nagata. 1988. Distribution of gorse (Ulex europaeus L.) a noxious weed in Hawaii. <u>Hawaiian Botanical Society Newsletter</u> 27(3): 110-117.
- Marler, M. J., C. A. Zabinski and R. M. Callaway. 1999. Mycorrhizae indirectly enhance competitive effects of an invasive forb on a native bunchgrass. <u>Ecology</u> 80(4): 1180-1186.
- Massey, J. E. 1978. Lake Waiau: a study of a tropical alpine lake, past and present. Honolulu, HI, University of Hawaii at Manoa: 130 p.
- Massey, J. E. 1979. The diatoms of contemporary and ancient sediments from Lake Waiau, Hawaii, and their geochemical environment. Review of Palaeobotany and Palynology 27(1): 77-83.
- Matthes, U., B. D. Ryan and D. W. Larson. 2000. Community structure of epilithic lichens on the cliffs of the Niagara Escarpment, Ontario, Canada. Plant Ecology 148: 233-244.
- Mauna Kea Management Board. 2004. Minutes Regular Meeting January 27, 2004.
- Mauna Kea Management Board. 2007. Minutes Regular Meeting March 6, 2007.
- McCoy, P. C. 1977. The Mauna Kea Adze Quarry Project: a summary of the 1975 field investigations. Journal of the Polynesian Society **86**(2): 223-244.
- McCoy, P. C. 1984. Mauna Kea summit region survey: a summary of the 1984 fieldwork. Honolulu, HI, Dept. of Anthropology, Bishop Museum: 11 p.
- McCoy, P. C. and R. A. Gould. 1977. Alpine archaeology in Hawaii. Archaeology 30(4): 234-243.
- McCoy, P. C., S. Collins, et al. 2009. Draft Report: A Cultural Resources Management Plan for the University of Hawaii Management Areas on Mauna Kea, Ka'ohe Ahupua'a, Hāmākua District, Hawai'i Island, State of Hawaii. TMK: (3) 4-4-015: 09, 12 2, 015; A Sub-Plan for the Mauna Kea Comprehensive Management Plan, Prepared by Pacific Consulting Services, Inc. for Office of Mauna Kea Management.
- McDonald, T. L. and P. H. Geissler. 2004. Systematic and Stratified Sampling Designs In Long-Term Ecological Monitoring Studies. West Inc. and U.S. Geological Survey. Retrieved from http://science.nature.nps.gov/im/monitor/docs/SampleDesigns.doc.
- McDonnell, M. J. and E. W. Stiles. 1983. The structural complexity of old field vegetation and the recruitment of bird-dispersed plant species. <u>Oecologia</u> 56: 109-116.

- McIntosh, R. W. and D. N. Milstein. 1964. The study of the Pohakuloa-Mauna Kea recreation area, County of Hawaii, State of Hawaii. <u>Mauna Kea</u>. Honolulu, HI, Bureau of Business Research, College of Business Administration, University of Hawaii.
- McLoud, P. 2003. "A Day in the Life of a Mauna Kea Ranger." from http://www.malamamaunakea.org/management.php?article_id=48.
- MCM Planning. 1988. Final supplemental environmental impact statement VLBA Antenna Facility, Mauna Kea, Hamakua, Hawaii, September, 1988. Amendment to the Mauna Kea Science Reserve Complex Development Plan. Prepared for The National Radio Astronomy Observatory, Socorro, New Mexico.
- MCM Planning. 1994. Project description and environmental review, Smithsonian Astrophysical Observatory Submillimeter Array Telescope, Mauna Kea, Hamakua Hawaii. Prepared for Smithsonian Astrophysical Observatory, Cambridge, Massachusetts, January 1994.
- McNarie, A. D. 2004. Mercury on the mountain. Hawaii Island Journal Volume, DOI:
- Medeiros, M. J. 2008. Personal communication from Matthew Medeiros, U.C. Berkeley, to Jennifer Garrison, SRGII, regarding moth species diversity on Mauna Kea.
- Meinecke, W. H. 1916. Insects from the crater of Mauna Loa. <u>Proceedings of the Hawaiian Entomological Society</u> 3(4): 285-286.
- Mellow, J. 2008. Personal communication from J. Mello, Biologist, Hawaii Division of Forestry and Wildlife to Colleen Bergmanis SRGII.
- Melvin, D. 1988. Poliahu: snow goddess of Mauna Kea. Spirit of Aloha 13(6): 51.
- Menard, T. 2001. Activity patterns of the Hawaiian hoary bat (Lasiurus cinereus semotus) in relation to reproductive time periods. <u>Zoology Department</u>. Honolulu, HI, University of Hawaii: 149 p.
- Menard, T. 2008. Personal communication from Theresa Menard, Hawaiian hoary bat expert, to Jennifer Garrison, SRGII, regarding habitat use and status of Hawaiian hoary bat on Mauna Kea.
- Menviel, L. (n.d.). "Study of Lake Waiau (Big Island of Hawaii) sediments as a possible paleoclimatologic record. Identification and isolation of diatoms from Lake Waiau sediments." from http://www.soest.hawaii.edu/oceanography/students/lmenviel/lakewaiau.html.
- Mesick, L. S. 1909. Hawaii Redivivus. Paradise of the Pacific 22(9): 21-24.
- Meyer, W. M. 2008. Personal communication from Marty Meyer, University of Hawai'i, to Jennifer Garrison, SRGII, regarding the garlic snail, *Oxychilus alliarius*.
- Meyer, W. M., III. 2005. Potential impact of an abundant introduced omnivorous land snail, Oxychilus alliarius on native land snails on the Island of Hawaii. <u>Tentacle</u> 13: 16-17.
- Meyer, W. M., III. 2006. Helicinids and endodontids alive in Hawaii! A comment on the conservation of family level land snail diversity in Hawaii. Tentacle 14: 29.
- Mitchell, C., C. Ogura, et al. 2005a. Hawaii's comprehensive wildlife conservation strategy. Honolulu, HI, As submitted to the National Advisory Acceptance Team, Dept. of Land and Natural Resources.
- Mitchell, C., C. Ogura, D. Meadows, A. Kane, L. Strommer, S. Fretz, D. Leonard and A. McClung. 2005b. 'Io, *Buteo solitarius*. <u>Hawaii's comprehensive wildlife conservation strategy</u>. Honolulu, HI, As submitted to the National Advisory Acceptance Team, Dept. of Land and Natural Resources.
- Miyashita, A., N. Takato, et al. 2004. Statistics of weather data, environmental data and the seeing of the Subaru Telescope. Ground-based Telescopes. J. M. Oschmann, Proc. of SPIE. **5489**: 207-217.
- MKMB Environment Committee. 2007. Personal communication from MKMB Environment Committee, to SRGII, regarding road grading at Mauna Kea.
- Montgomery, M. 2008. Personal Communication from Michelle Montgomery, USGS, to Jennifer Garrison, SRGII, about terrestrial arthropod monitoring techniques.
- Montgomery, S. L. 1988. A report on the invertebrate fauna found on the proposed NRAO VLBA antenna facility site, Mauna Kea Science Reserve, Mauna Kea, Hamakua, Hawaii. Appendix G in: MCM Planning. 1988. Final supplemental environmental impact statement VLBA Antenna Facility, Mauna Kea, Hamakua, Hawaii, September, 1988. Amendment to the Mauna Kea Science Reserve

- Complex Development Plan. Prepared for The National Radio Astronomy Observatory, Socorro, New Mexico.
- Montgomery, S. L. and F. G. Howarth. 1980. Records of mummified leiothrix from the summits of Mauna Loa and Mauna Kea. <u>Elepaio</u> 41(4): 30-31.
- Moore, J. G. 2001. Density of basalt core from Hilo drill hole, Hawaii. <u>Journal of Volcanology and</u> Geothermal Research v. 112(no. 1-4): p. 221-230.
- Moore, J. G. and D. A. Clague 1992. Volcano growth and evolution of the island of Hawaii. <u>Geological Society of America Bulletin</u> **104**(10): 1471-1484.
- Moore, J. G., D. A. Clague, et al. 1990. Subsidence and volcanism of the Haleakala Ridge, Hawaii. Journal of Volcanology and Geothermal Research v. 42: p. 273-284.
- Moore, J. G., S. C. Porter, et al. 1993. Glaciation of Haleakala Volcano, Hawaii [abs.]. <u>Geological Society of America Abstracts with Programs</u> v. 25(no. 5): p. 123-124.
- Moulton, M. P., K. E. Miller and E. A. Tillman. 2001. Patterns of success among introduced birds in the Hawaiian Islands. In: Evolution, ecology, conservation, and management of Hawaiian birds: A vanishing avifauna. <u>Studies in Avian Biology No. 22.</u> J. M. Scott, S. Conant and C. van Riper, III. Lawrence, KS, Cooper Ornithological Society: 31-46.
- Mountainspring, S. and J. M. Scott. 1985. Interspecific competition among Hawaiian forest birds. <u>Ecological Monographs</u> 55(2): 219-239.
- Mueller-Dombois, D. and F. R. Fosberg. 1998. *Vegetation of the Tropical Pacific Islands*. New York, Springer-Verlag.
- Mueller-Dombois, D. and V. J. Krajina 1968. Comparison of east-flank vegetations on Mauna Loa and Mauna Kea, Hawaii. <u>Proceedings of the Symposium on Recent Advances in Tropical Ecology</u>. Varanasi, India., Varanasi International Society for Tropical Ecology: p. 508-520.
- Mull, M. E. 1979. Palila triumphs in court. Elepaio 40(1): 4.
- Mull, M. E. 1977. Palila at Hale Pohaku. Elepaio 38(1): 10.
- Munro, G. C. 1945. Pekin nightingale on mountain top. <u>Elepaio</u> 6(3): 19-20.
- Nagata, S. 2007. Personal communication from Stephanie Nagata, OMKM, to Kristin Duin and Jennifer Garrison, SRGII, regarding MKSR.
- Nahakuelua, R. 2008. Personal communication from R. Nahakuelua, MKOSS to Colleen Bergmanis, SRGII.
- NASA. 2005. Final environmental impact statement for the Outrigger Telescopes Project: Mauna Kea Science Reserve, Island of Hawai'i. National Aeronautics and Space Administration, Universe Division, Science Mission Directorate Washington, D.C.
- National Invasive Species Council. 2008. 2008-2012 National Invasive Species Management Plan. Washington DC, National Invasive Species Council, Department of the Interior: 35 pp.
- National Park Service. 2006. Guidance for Designing an Integrated Monitoring Program. from http://science.nature.nps.gov/im/monitor/GoalsObjectives.cfm.
- Neal, M. C. 1939. A list of mosses and vascular plants collected on Mauna Kea, August 1935. Proceedings of the Hawaiian Academy of Science **34**: 13.
- Neal, M. C. 1939. Vegetation of Lake Waiau, Hawaii. Paradise of the Pacific 51(10): 7, 32.
- Newton, G. A. and V. P. Claassen. 2003. <u>Rehabilitation of disturbed lands in California: a manual for decision-making. Special publication 123</u>. Sacramento, CA, California Department of Conservation, California Geological Survey.
- Noguchi, Y., H. Tabuchi, et al. 1987. Physical factors controlling the formation of patterned ground on Haleakala, Maui. Geografiska Annaler, Series A, Physical Geography **69**(2): 329-342.
- NPS. 1994. Mauna Kea, U.S. Department of the Interior, National Park Service, National Natural Landmarks Progam: 36 p.
- Nullet, D. and T. W. Giambelluca. 1990. Winter evaporation on a mountain slope, Hawaii. <u>Journal of Hydrology</u> **112**: 257-265.

- Nullet, D., J. O. Juvik, et al. 1995. A Hawaiian mountain climate cross-section. <u>Climate Research</u> 5: 131-137.
- Oakley, K. L., L. P. Thomas and S. G. Fancy. 2003. Guidelines for long-term monitoring protocols. Wildlife Society Bulletin 31(4): 1000-1003.
- Oboyski, P. T. 2008. Personal communication from Peter Oboyski to Jennifer Garrison, SRGII, regarding arthropods at Hale Pohaku and in the Mauna Kea Science Reserve.
- Oboyski, P. T., J. W. Slotterback and P. C. Banko. 2004. Differential parasitism of seed-feeding Cydia (Lepidoptera: Tortricidae) by native and alien wasp species relative to elevation in subalpine Sophora (Fabaceae) forests on Mauna Kea, Hawaii. <u>Journal of Insect Conservation</u> 8(2-3): 229-240.
- Odum, S. 1965. Germination of ancient seeds. <u>Dansk Botanisk Arkiv</u> 24: 70.
- Office of the Legislative Auditor. 1998. Audit of the management of Mauna Kea and the Mauna Kea Science Reserve: a report to the Governor and the Legislature of the State of Hawaii. Honolulu, The Auditor: v, 54 p.
- Office of the Legislative Auditor. 2005. Follow-up audit of the management of Mauna Kea and the Mauna Kea Science Reserve: a report to the Governor and the Legislature of the State of Hawaii. Honolulu, HI, Hawaii Legislature Office of the Legislative Auditor: 52 p.
- Ohio Agriculture Research and Development Center. n.d. Common cat's ear, Hypochoeris radicata. Ohio Perennial and Biennial Weed Guide.
- Okinaka, A. M. 2004. Results from a survey on managing access to the Mauna Kea scientific reserve., Prepared for the Office of Mauna Kea Management by Alton M. Okinaka, July 19, 2004.
- Olson, S. L. and H. F. James. 1982. Fossil birds from the Hawaiian Islands: evidence for wholesale extinction by man before western contact. <u>Science</u> 217: 633-635.
- OMKM Rangers. 2007. Personal communication from OMKM Rangers (Pablo McLoud and Shane) regarding Ranger duties and visitors to Mauna Kea. Kristin Duin SRGII.
- Pacific Analytics. 2000. Wekiu Bug Mitigation Report. Prepared for the Keck Outrigger Telescopes Project, Jet Propulsion Laboratory, California Institute of Technology. Pacific Analytics LLC Albany, OR.
- Pacific Analytics. 2001a. Wekiu Bug Mitigation Plan. Appendix D in Environmental assessment for the Outrigger Telescopes Project: Mauna Kea Science Reserve, Island of Hawai°i. National Aeronautics and Space Administration, Office of Space Science.
- Pacific Analytics. 2001b. Wekiu Bug Monitoring Plan. Prepared for the Outrigger Telescopes Project, Jet Propulsion Laboratory, California Institute of Technology. Albany, OR.
- Pacific Analytics. 2004. Botanical survey of the Hale Pohaku mid-elevation facilities construction staging area. Prepared for The Outrigger Telescopes Project, National Aeronautics and Space Administration by Pacific Analytics LLC Albany, OR.
- PBS Hawaii. 2004. First Light.
- Pejchar, L. and J. Jeffrey. 2004. Sap-feeding behavior and tree selection in the endangered akiapolaau (Hemignathus munroi) in Hawaii. <u>Auk</u> 121(2): 548-556.
- Pejchar, L., K. D. Holl and J. L. Lockwood. 2005. Hawaiian honeycreeper home range size varies with habitat: implications for native acacia koa forestry. <u>Ecological Applications</u> 15(3): 1053-1061.
- Peng, L. and J. W. King 1992. A late quaternary geomagnetic secular variation record from Lake Waiau, Hawaii, and the question of the Pacific nondipole low. <u>Journal of Geophysical Research</u> v. 97(no. B4): p. 4407-4424.
- Polhemus, D. A. 1998. Nysius aa (Heteroptera: Lygaeidae), a new species of micropterous wekiu bug from the summit of Mauna Loa volcano, Hawaii. <u>Proceedings of the Entomological Society of Washington 100(1): 25-31.</u>
- Polhemus, D. A. 2001. A preliminary survey of Wekiu Bug populations at Puu Hau Kea, in the Mauna Kea Ice Age Natural Area Reserve, Hawaii Island, Hawaii. Smithsonian Institution Washington, D.C.

- Polhemus, D. A. 2004. Further studies on the genus Orthotylus (Heteroptera: Miridae) in the Hawaiian Islands, with descriptions of thirty-four new species. J. New York Entomol. Soc. 112(4): 227–333.
- Porter, S. C. 1972a. Buried caldera of Mauna Kea Volcano, Hawaii. Science 175: 1458-1460.
- Porter, S. C. 1972b. Distribution, morphology, and size frequency of cinder cones on Mauna Kea Volcano, Hawaii. <u>Geological Society of America Bulletin</u> **83**(12): 3607-3612.
- Porter, S. C. 1973a. <u>Late Quaternary glaciations and tephrochronology of Mauna Kea, Hawaii [abs.]</u>. International Union for Quaternary Research Program, 9th Congress, Proceedings.
- Porter, S. C. 1973b. Stratigraphy and chronology of late quaternary tephra along the South Rift Zone of Mauna Kea Volcano, Hawaii. <u>Geological Society of America Bulletin</u> **84**(6): 1923-1940.
- Porter, S. C. 1975. Late quaternary glaciation and tephrochronology of Mauna Kea, Hawaii. <u>Quaternary Studies</u> 13: 247-251.
- Porter, S. C. 1979a. Hawaiian glacial ages. Quaternary Research 12: 161-187.
- Porter, S. C. 1979b. Quaternary stratigraphy and chronology of Mauna Kea, Hawaii: A 380,000-yr record of mid-Pacific volcanism and ice-cap glaciation. <u>Geological Society of America Bulletin, Part II</u> **90**(7): 908-1093.
- Porter, S. C. 1979c. Quaternary stratigraphy and chronology of Mauna Kea, Hawaii: a 380,000-yr record of mid-Pacific volcanism and ice-cap glaciation: Summary. <u>Geological Society of America Bulletin</u>, part 1 **90**: 609-611.
- Porter, S. C. 1986. Glaciation of Mauna Kea, Hawaii. Quaternary Science Reviews v. 5: p. 181-182.
- Porter, S. C. 1987. Pleistocene subglacial eruptions on Mauna Kea. <u>U.S. Geological Survey Professional Paper 1350</u>. R. W. Decker, T. L. Wright and P. H. Stauffer. **v. 1:** p. 587-598.
- Porter, S. C. 1997a. Late Pleistocene eolian sediments related to pyroclastic eruptions of Mauna Kea volcano, Hawaii. <u>Quaternary Research</u> **47**(3): 261-276.
- Porter, S. C. 1997b. Volcanic loess and dune sand related to late Pleistocene eruptions of Mauna Kea Volcano, Hawaii [abs.]. Geological Society of America Abstracts with Programs v. 29(no. 5): p. 57.
- Porter, S. C. 2005. Pleistocene snowlines and glaciation of the Hawaiian Islands. <u>Quaternary International</u> 138-139: 118-128.
- Porter, S. C. and R. A. Englund. 2006. Possible geologic factors influencing the distribution of the Wekiu Bug on Mauna Kea, Hawaii. Hawaii Biological Survey Report 2006-031. Prepared for the Office of Mauna Kea Management by S.C. Porter and R.A. Englund, Hawaii Biological Survey, Bishop Museum Honolulu, HI.
- Porter, S., M. Stuiver, et al. 1977. Chronology of Hawaiian glaciation. Science 195: 61-63.
- Pounds, J. A., M. P. L. Fogden and J. H. Campbell. 1999. Biological response to climate change on a tropical mountain. Nature 398: 611-615.
- Powers, W. E. and C. K. Wentworth. 1941. Air movements and weather in the Mauna Kea--Mauna Loa Saddle, Hawaii. <u>Bulletin of the American Meteorological Society</u> **22**: 6-13.
- Pratt, H. D. 1980. Intra-island variation in the elepaio on the island of Hawaii. Condor 82(4): 449-458.
- Pratt, T. K. 2008. Personal communication from Thane Pratt, USGS, to Jennifer Garrison, SRGII, regarding native birds at Hale Pohaku.
- Pratt, T. K., P. C. Banko, S. G. Fancy, G. D. Lindsey and J. D. Jacobi. 1997. Status and management of the Palila, an endangered Hawaiian honeycreeper, 1987-1996. <u>Pacific Conservation Biology</u> 3(4): 330-340.
- Preston, D. J. 2008. Personal communication from David Preston, Bishop Museum, to Jennifer Garrison, SRGII, regarding Wēkiu bug sampling locations.
- Raine, C. T. 1939. Meteorological reports of the Mauna Kea Expedition, 1935 (II); the meteorological observations at Lake Waiau, August 8-19, 1935. <u>Bulletin of the American Meteorological Society</u> **20**: 97-103.
- Ramsdale, A. S. 2004. Records of Coleoptera from the Alpine Zone of Mauna Kea, Hawaii. Bishop Museum Honolulu, HI.

- Richardson, M. 2002. Life in Mauna Kea's alpine desert. Endangered Species Bulletin 27(2): 22-23.
- Robichaux, R. H., E. A. Friar and D. W. Mount. 1997. Molecular genetic consequences of a population bottleneck associated with reintroduction of the Mauna Kea silversword (Argyoxiphium sandwicense ssp. Sandwicense (Asteraceae). Conservation Biology 11(5): 1140-1146.
- Robichaux, R. H., S. Bergfeld, M. Bruegmann, J. Canfield, P. Moriyasu, T. Rubenstein and T. W. Tunison, Frederick. 2000. Reintroducing Hawaii's silverswords. <u>Endangered Species Bulletin</u> 25(3): 22-23.
- Rock, J. F. C. 1913. The indigenous trees of the Hawaiian Islands. Honolulu, Published under patronage.
- Roderick, G. K. and R. G. Gillespie. 1998. Speciation and phylogeography of Hawaiian terrestrial arthropods. Molecular Ecology 7: 519-531.
- Rohr, J. R., C. G. Mahan and K. C. Kim. 2007. Developing a monitoring program for invertebrates: guidelines and a case study. Conservation Biology 21(2): 422-433.
- Rohrer, J. 2008. Personal communication from Joby Rohrer, Natural Resource Management Coordinator, US Army, to Jennifer Garrison, SRGII, regarding small mammal monitoring techniques.
- Rowland, S. K. and G. P. Walker 1990. Pahoehoe and aa in Hawaii: volumetric flow rate controls the lava structure. <u>Bulletin of Volcanology</u> **v. 52**: p. 615-628.
- Rundell, R. J. 2008. Personal communication from Rebecca Rundell, U. of Chicago, to Jennifer Garrison, SRGII, regarding snail survey and monitoring techniques.
- Rundell, R. J. and R. H. Cowie. 2003. Growth and reproduction in Hawaiian succineid land snails <u>Journal</u> of Molluscan Studies 69: 288–289.
- Sato, H. H., W. Ikeda, et al. 1973. Soil Survey of the Island of Hawaii, State of Hawaii. United States Department of Agriculture Soil Conservation Service.
- Schemnitz, S. D., Ed. 1980. Wildlife management techniques manual. Washington D.C., The Wildlife Society.
- Schwartz, C. W. and E. R. Schwartz. 1951. An ecological reconnaissance of the pheasants in Hawaii. <u>Auk</u> 68(3): 281-314.
- Scott, J. M., S. Mountainspring, C. van Riper, III, C. B. Kepler, J. D. Jacobi, T. A. Burr and J. G. Giffin. 1984. Annual variation in the distribution, abundance, and habitat response of the palila (Loxioides bailleui). <u>Auk</u> 101(4): 647-664.
- Scott, J. M., S. Mountainspring, F. L. Ramsey and C. B. Kepler. 1986. Forest bird communities of the Hawaiian Islands: their dynamics, ecology, and conservation. Los Angeles, CA, Cooper Ornithological Society.
- Scowcroft, P. G. 1983. Tree cover changes in Mamane (Sophora chrysophylla) forests grazed by sheep and cattle. Pacific Science 37(2): 109-119.
- Scowcroft, P. G. and C. E. Conrad. 1992. Alien and native plant response to release from feral sheep browsing on Mauna Kea. <u>Alien Plant Invasions in Native Ecosystems of Hawaii: Management and Research C. P. Stone, C. W. Smith and J. T. Tunison. Honolulu, University of Hawaii Cooperative National Park Resources Studies Unit pp 625-665.</u>
- Scowcroft, P. G. and H. F. Sakai. 1983. Impact of feral herbivores on mamane forests of Mauna Kea, Hawaii: bark stripping and diameter class structure. Journal of Range Management 36(4): 495-498.
- Scowcroft, P. G. and J. G. Giffin. 1983. Feral herbivores suppress mamane and other browse species on Mauna Kea, Hawaii. <u>Journal of Range Management</u> 36(5): 638-645.
- Settle, M. 1979. The structure and emplacement of cinder cone fields. <u>American Journal of Science</u> **279**(10): 1089-1107.
- Sharifi, M. R., A. C. Gibson and P. W. Rundel. 1997. Surface dust impacts on gas exchange in Mojave Desert shrubs. Journal of Applied Ecology 34: 837-846.
- Sharp, W. D. and P. R. Renne 2005. The 40 Ar/39 Ar dating of core recovered by the Hawaii Scientific Drilling Project (phase 2), Hilo, Hawaii. G³ (Geochemistry, Geophysics, Geosystems) v. 6(no. Q04G17): doi:10.1029/2004GC000846.

- Shaw, R. B. 1997. Rare plants of the Pohakuloa Training Area, Hawaii. Retrieved Sept. 23, 2007, from https://www.denix.osd.mil/denix/Public/ES-
 - Programs/Conservation/Legacy/HawaiiPlants/hiplant.html#asfr.
- Sheldon, N. D. 2006. Quaternary Glacial-Interglacial Climate Cycles in Hawaii. <u>Journal of Geology</u> **114**(3): 367-376.
- Sherrod, D. R., J. M. Sinton, et al. 2007. Geologic map of the State of Hawai'i, U.S. Geological Survey: 83 p.
- Skottsberg, C. J. F. 1931. Remarks on the flora of the high Hawaiian volcanoes. <u>Acta Horti Gothoburgensis</u> 6: 45-65.
- Smith, C. 2003. Petition to list the Wekiu bug (Nysius Wekiuicola) as an endangered species with critical habitat designation. Submitted to the United States Department of the Interior, Fish and Wildlife Service, by KAHEA: the Hawaiian Environmental Alliance, May 22, 2003.
- Smith, C. W. 1985. Impact of alien plants on Hawai'i's native biota. <u>Hawaii's terrestrial ecosystems:</u> <u>preservation and management</u>. C. P. Stone and M. J. Scott. Honolulu, HI, Cooperative National Park Resources Studies Unit, University of Hawaii: 180-250.
- Smith, C. W. and J. T. Tunison 1992. Fire and alien plants in Hawai'i: Research and management implications for native ecosystems. <u>Alien Plant Invasions in Native Ecosystems of Hawai'i: Management and Research C. P. Stone, C. W. Smith and J. T. Tunison. Honolulu, University of Hawaii Cooperative National Park Resources Studies Unit pp 394-408.</u>
- Smith, C. W., W. J. Hoe and P. J. O'Conner. 1982. Botanical Survey of the Mauna Kea summit above 13,000 feet. Prepared for Group 70. October 1982.
- Smith, D. R. 1967. New Localities for Hawaiian Mosses. The Bryologist 70(2): 237-245.
- Smith, R. S., R. S. Shiel, et al. 2008. Long-term change in vegetation and soil microbial communities during the phased restoration of traditional meadow grassland. <u>Journal of Applied Ecology</u> **45**: 670–679.
- Snetsinger, T. J. 1995. Observations of Pueo nests on the slopes of Mauna Kea. <u>Elepaio</u> 55(1): 3.
- Snetsinger, T. J., S. G. Fancy, J. C. Simon and J. D. Jacobi. 1994. Diets of owls and feral cats in Hawaii. <u>Elepaio</u> 54(8): 47-50.
- Society for Ecological Restoration International Science & Policy Working Group 2004. The SER International Primer on Ecological Restoration. Tucson, AZ, Society for Ecological Restoration International: 13 p.
- Sridhar, K., M. L. Jackson, et al. 1978. Oxygen isotopic ratios of quartz from wind-erosive soils of southwestern United States in relation to aerosol dust. <u>Soil Science Society of America Journal</u> **42**(1): 158-162.
- Stampe, E. D. and C. C. Daehler. 2003. Mycorrhizal species identity affects plant community structure and invasion: a microcosm study. Oikos 100: 362-372.
- Stearns, H. T. 1945. Glaciation of Mauna Kea, Hawaii. <u>Geological Society of America Bulletin</u> **56**(3): 267-274.
- Stearns, H. T. 1946. Geology of Mauna Kea: previous investigations. <u>Geology and ground-water resources of the island of Hawaii</u>. H. T. Stearns and G. A. MacDonald. Honolulu, HI, Dept. of Public Lands, Division of Hydrography: p. 150-169.
- Stearns, H. T. and G. A. Macdonald. 1946. Geology and Ground-Water Resources of the Island of Hawaii, Hawaii Division of Hydrography. **Bulletin 9**.
- Stemmerman Kjargaard, M. 1985. Avian survey of the Hale Pohaku area. Appendix B in: MCM Planning and University of Hawaii. 1985. Final supplemental environmental impact statement for construction camp housing, amendment to the Mauna Kea Science Reserve Complex Development Plan. Hale Pohaku, Hamakua, Mauna Kea, Hawaii. Prepared for MCM Planning Volcano, HI.

- Stemmerman, M. 1979. Avian Survey. Appendix in: Group 70. 1980. Revised environmental impact statement for Hale Pohaku: mid-elevation facilities master plan, Hamakua, Mauna Kea, Hawaii. Prepared for Group 70 Honolulu, HI.
- Stiles, E. W. and D. W. White. 1986. Seed deposition patterns: influence of season, nutrients, and vegetation structure. <u>Frugivores and seed dispersal</u>. A. Estrada and T. H. Fleming. Dordrecht, Dr W. Junk Publishers: 45-54.
- Still, C. J., P. N. Foster and S. H. Schneider. 1999. Simulating the effects of climate change on tropical montane cloud forests. <u>Nature</u> 398: 608-610.
- Stinson, K. A., S. A. Campbell, J. R. Powell, B. E. Wolfe, R. M. Callaway, G. C. Thelen, S. G. Hallett, D. Prati and J. N. Klironomos. 2006. Invasive plant suppresses the growth of native tree seedlings by disrupting belowground mutualisms. <u>PLoS Biology</u> 4(5): e140.
- Stone, C. P. 1985. Alien animals in Hawaii's native ecosystems: towards controlling the adverse effects of introduced vertebrates. <u>Hawaii's terrestrial ecosystems: preservation and management</u>. C. P. Stone and J. M. Scott. Honolulu, Uhiv. Hawaii Press: 251-297.
- Stone, C. P. and S. J. Anderson. 1988. Introduced Animals in Hawaii's Natural Areas. Proceedings of the Thirteenth Vertebrate Pest Conference (1988). Vertebrate Pest Conference Proceedings Collection. University of Nebraska, Lincoln. Retrieved from http://digitalcommons.unl.edu/vpcthirteen/28/.
- Stopler, E. M., D. J. DePaolo, et al. 1996. Introduction to special section: Hawaii Scientific Drilling Project. <u>Journal of Geophysical Research</u> **101**(B5): 11,593-11,598.
- Swezey, O. H. and F. X. Williams. 1932. Insects from the summit of Mauna Kea. <u>Proceedings of the Hawaiian Entomological Society</u> 8(1): 191-192.
- Thomas, D. 2008. Personal communication from D. Thomas, MKMB Environment Committee Member to SRGII.
- Three Mountain Alliance 2007. Three Mountain Alliance Management Plan: 82 p.
- Toma, J. 2008. Personal communication from J. Toma, Environmental Health Specialist, State Department of Health Indoor and Radiological Health (IRH) Branch regarding noise monitoring at Mauna Kea. C. B. SRGII.
- Tomlinson, A. 2007. Personal communication from Andrew Tomlinson, PCSI, to Jennifer Garrison, SRGII, regarding location of Mauna Kea silversword in the MKSR.
- Traveset, A. and D. M. Richardson. 2006. Biological invasions as disruptors of plant reproductive mutualisms. Trends in Ecology and Evolution 21(4): 208-216.
- Tunison, J. T. 1992. Fountain grass control in Hawai'i Volcanoes National Park: Management considerations and strategies. <u>Alien Plant Invasions in Native Ecosystems of Hawai'i: Management and Research C. P. Stone, C. W. Smith and J. T. Tunison. Honolulu, University of Hawaii Cooperative National Park Resources Studies Unit pp 376-393.</u>
- U. S. Geological Survey. 1997a. "Earthquake Hazards." from http://pubs.usgs.gov/gip/hazards/earthquakes.html.
- U. S. Geological Survey. 1997b. "Lava Flow Hazard Zone Maps: Mauna Kea and Kohala." from http://pubs.usgs.gov/gip/hazards/maunakea-kohala.html.
- U. S. Geological Survey. 2008a. "Halema`uma`u gas plume becomes ash-laden." USGS HVO News Release, from http://hvo.wr.usgs.gov/pressreleases/pr03 24 08.html.
- U. S. Geological Survey. 2008b. "New gas vent in Halema'uma'u crater doubles sulfur dioxide emission rates" USGS HVO News Release, from http://hvo.wr.usgs.gov/pressreleases/pr03_14_08.html.
- U.S. Geological Survey. 2002, May 22, 2002. "Mauna Kea: Hawai`i's Tallest Volcano." from http://hvo.wr.usgs.gov/volcanoes/maunakea/.
- Ueno, M. 1936. Cladocera of Mauna Kea, Hawaii. <u>Bernice P. Bishop Museum Occasional Papers</u> **12**(11): 9 p.
- Ugolini, F. C. 1974. Hydrothermal origin of the clays from the upper slopes of Mauna Kea, Hawaii. <u>Clay and Clay Minerals</u> **22**: 189-194.

- University of Hawaii. 1985. Conservation District Use Application for the University of Hawaii's Mauna Kea Management Plan. Mauna Kea. Honolulu, HI, Submitted to the State of Hawaii, the Board, Dept. of Land and Natural Resources. February 22, 1985.
- University of Hawaii. 2000. Voices and visions of Mauna Kea: Mauna Kea Science Reserve master plan and implementation process summary. March 2000. <u>Master Plan</u>. Honolulu, HI, University of Hawaii: 7 p.
- University of Hawaii Institute for Astronomy. 1981. Program summary and research development plan for the Mauna Kea Science Reserve and related facilities. <u>Master Plan</u>. Honolulu, HI, Institute for Astronomy, University of Hawaii at Manoa: 49 p.
- University of Hawaii Institute for Astronomy. 2002. Final environmental assessment for the W. M. Keck Observatory Outrigger Telescopes Project. March 2002. <u>Outrigger</u>. Hilo, HI.
- US Forest Service Institute of Pacific Islands Forestry. 2005. "Pacific Island Ecosystems at Risk (PIER) Risk Assessments." from http://www.hear.org/pier/wralist.htm.
- USFWS. 1977. Determination of critical habitat for six endangered species. <u>Federal Register</u> 42(155): 40685-40690.
- USFWS. 1994. Recovery plan for the Mauna Kea Silversword (Argyroxiphium sandwicense ssp. sandwicense). from http://ecos.fws.gov/docs/recovery_plans/1993/930930a.pdf.
- USFWS. 1998a. Recovery plan for the Hawaiian hoary bat (Lasiurus cinereus semotus). U.S. Fish and Wildlife Service Portland, OR. Retrieved from http://ecos.fws.gov/docs/recovery_plans/1998/980511b.pdf.
- USFWS. 1998b. The Final Recovery Plan for Four Species of Hawaiian Fern. Pacific Islands Endangered Species (web page) Retrieved Sept. 23, 2007, from http://www.fws.gov/pacificislands/recoveryplans.html.
- USFWS. 2002. Endangered and Threatened wildlife and plants; Designations of critical habitat for plant species from the Island of Hawaii, Hawaii. Federal Register 67(102): 36967-37016.
- USFWS. 2004. Draft revised recovery plan for the Nene or Hawaiian goose (*Branta sandvicensis*). U.S. Fish and Wildlife Service Portland, OR.
- USFWS. 2006. 50 CFR Part 17: Endangered and Threatened Wildlife and Plants; Review of Native Species That Are Candidates or Proposed for Listing as Endangered or Threatened; Annual Notice of Findings on Resubmitted Petitions; Annual Description of Progress on Listing Actions. Volume 71, Number 176: 53755-53835. Retrieved from http://www.fws.gov/policy/library/06-7375.html.
- Valier, K. 1995. Ferns of Hawai'i. Honolulu, HI, University of Hawai'i Press.
- Van de Ven, C. M., S. B. Weiss and W. G. Ernst. 2007. Plant species distributions under present conditions and forecasted for warmer climates in an arid mountain range. <u>Earth Interactions</u> 11(9): 1-33.
- van Riper, C., III, J. M. Scott and D. M. Woodside. 1978. Distribution and abundance patterns of the palila on Mauna Kea, Hawaii. <u>Auk</u> 95(3): 518-527.
- van Riper, C., III, S. G. van Riper and W. R. Hansen. 2002. Epizootiology and effect of avian pox on Hawaiian forest birds. Auk 119(4): 929-942.
- van Riper, C., III, S. G. van Riper, M. L. Goff and M. Laird. 1986. The epizootiology and ecological significance of malaria in Hawaiian land birds. <u>Ecological Monographs</u> 56(4): 327-344.
- van Riper, C., III. 1976. Aspects of house finch breeding biology. Condor 78(2): 224-229.
- van Riper, C., III. 1978. Discovery of the yellow-fronted canary on Mauna Kea, Hawaii. <u>'Elepaio</u> 38(9): 99-100.
- van Riper, C., III. 1980a. Observations on the breeding of the palila Psittirostra bailleui of Hawaii. <u>Ibis</u> 122: 462-475.
- van Riper, C., III. 1980b. The phenology of the dryland forest of Mauna Kea, Hawaii, and the impact of recent environmental perturbations. <u>Biotropica</u> 12(4): 282-291.

- van Riper, C., III. 1984. The influence of nectar resources on nesting success and movement patterns of the common Amakihi (Hemignathus virens). <u>Auk</u> 101(1): 38-46.
- van Riper, C., III. 1987. Breeding ecology of the Hawaii common amakihi. Condor 89(1): 85-102.
- van Riper, C., III. 1995. Ecology and breeding biology of the Hawaii elepaio (Chasiempis sandwichensis bryani). <u>Condor</u> 97(2): 512-527.
- VanderWerf, E. A. 2008. Personal communication from Eric VanderWerf to Jennifer Garrison, SRGII.
- VIS Staff. 2007. Personal communication from VIS staff regarding visitors to Mauna Kea. Kristin Duin SRGII.
- Vitousek, P. M. and L. R. Walker. 1989. Biological invasion by Myrica faya in Hawaii: plant demography, nitrogen fixation, ecosystem effects. <u>Ecological Monographs</u> 59(3): 247-265.
- Wada, L. 2008. Personal communication from Lorena Wada, US FWS, regarding wekiu bug (Nysius wekiuicola) Candidate Conservation Agreement. Jennifer Garrison SRGII.
- Wagner, W. L., D. R. Herbst and S. H. Sohmer. 1990. *Manual of the Flowering Plants of Hawai'i*. Honolulu, Bishop Museum Press.
- Wainscoat, R. J. 2007. Protection of Hawaii's Observatories from Light Pollution. <u>Starlight: A Common Heritage</u>. C. Marín and J. Jafari, Starlight Initiative: 379-384.
- Walker, D. A. and K. R. Everett. 1987. Road dust and its environmental impact on Alaskan taiga and tundra. <u>Arctic and Alpine Research</u> **19**(4): 479-489.
- Walker, G. P. L. 1990. Geology and volcanology of the Hawaiian Islands. <u>Pacific Science</u> v. 44(no. 4): p. 315-347.
- Walker, M. F. 1983. A comparison of observing conditions on the summit cones and shield of Mauna Kea. <u>Publications of the Astronomical Society of the Pacific</u> **95**: 903-918.
- Warner, R. E. 1960. A forest dies on Mauna Kea; how feral sheep are destroying an Hawaiian woodland. Pacific Discovery 13(2): 6-14.
- Washington, H. S. 1923. Petrology of the Hawaiian Islands; I. Kohala and Mauna Kea, Hawaii. <u>American</u> Journal of Science, 5th Series 5: 465-502.
- Washington, H. S. 1923. The density of the earth as calculated from the densities of Mauna Kea and Haleakala. Journal of the Washington Academy of Sciences **13**(21): 453-456.
- Weed Society of Queensland. 2005. Telegraph Weed. Retrieved November 23, 2007, from http://www.wsq.org.au/heterotheca.html.
- Weltzin, J. F., R. T. Belote and N. J. Sanders. 2003. Biological Invaders in a Greenhouse World: Will Elevated CO₂ Fuel Plant Invasions? <u>Frontiers in Ecology and the Environment</u> 1(3): 146-153.
- Wentworth, C. K. 1935. Mauna Kea, the white mountain of Hawaii. Mid-Pacific Magazine **48**(4): 290-296.
- Wentworth, C. K. 1939. Suncups in Hawaiian snowbanks. Paradise of the Pacific 51(12): 47-48.
- Wentworth, C. K. 1940. Ablation of snow under the vertical sun in Hawaii. <u>Amercican Journal of Science</u> **238**(2): 112-116.
- Wentworth, C. K. and W. E. Powers 1941. Multiple glaciation of Mauna Kea, Hawaii. <u>Geological Society</u> of America Bulletin **52**: 1193-1218.
- Wentworth, C. K. and W. E. Powers. 1943. Glacial springs on the Island of Hawaii. <u>Journal of Geology</u> 60: 542-547.
- Wentworth, C. K., J. W. Coulter, et al. 1935. <u>The Mauna Kea expedition of 1935: Hawaiian Academy of Science</u>. Honolulu, Hawaii., Bernice P. Bishop Museum.
- Werner, B. T. and B. Hallet 1993. Numerical simulation of self-organized stone stripes. <u>Nature</u> **361**(6408): 142-145.
- West, H. B., M. O. Garcia, et al. 1988. Nature and cause of compositional variation among the alkalic cap lavas of Mauna Kea Volcano, Hawaii. <u>Contributions to Mineralogy and Petrology</u> **100**(3): 383-397.
- Wetterer, J. K., P. C. Banko, L. P. Laniawe, J. W. Slotterback and G. J. Brenner. 1998. Nonindigenous ants at high elevations on Mauna Kea, Hawaii. <u>Pacific Science</u> 52(3): 228-236.

- White, T. A. and K. S. Hashisaki 1996. Taking a broader view: seeking common ground for the restoration paradigm. The role of restoration in ecosystem management. D. L. Pearson and C. V. Klimas. Madison, WI, Society for Ecological Restoration: 5-14.
- Williams, A. C. 1994. Effects of non-native grasses on a dry subalpine forest native to Hawaii and a discussion of effects of canopy on understory productivity, University of California at Berkeley: 46 pp.
- Williams, D. G. and R. A. Black. 1993. Phenotypic variation in contrasting temperature environments: Growth and photosynthesis in Pennisetum setaceum from different altitudes on Hawaii. <u>Functional Ecology</u> 7(5): 623-633.
- Williams, D. G., R. N. Mack and R. A. Black. 1995. Ecophysiology of introduced Pennisetum setaceum on Hawaii: The role of phenotypic plasticity. <u>Ecology</u> 76(5): 1569-1580.
- Wilson, C. 2008. Fighting invasive species in Hawaii. The Honolulu Advertiser January 14, 2008.
- Wilson, K. 2008. Personal communication from K. Wilson, Logistics Clerk, Mauna Kea Observatory Support Services to Colleen Bergmanis SRGII.
- Winterer, E. L., D. M. Hussong, et al. 1989. The Hawaiian-Emperor Chain. Boulder, CO, Geological Society of America. v. N: p. 187-287; pls. 1A, 1B, 2A, 2B, 3A, and 5 (folded maps) in separate box include Hawaii.
- Wittenberg, R. and M. J. W. Cock, Eds. 2001. <u>Invasive Alien Species: A Toolkit of Best Prevention and Management Practices</u>. Wallingford, Oxon, UK, CAB International.
- Wolfe, E. W. and J. Morris 1996a. Geologic map of the Island of Hawaii. <u>U.S. Geological Survey Miscellaneous Investigations Series I-2524-A.</u>
- Wolfe, E. W. and J. Morris, compilers 1996b. Sample data for the geologic map of the island of Hawaii. U.S. Geological Survey Miscellaneous Investigations Series Map I-2524-B: 3.
- Wolfe, E. W., W. S. Wise and G. B. Dalrymple. 1997. The geology and petrology of Mauna Kea volcano, Hawaii: a study of postshield volcanism. U.S. Geological Survey Professional Paper 1557. U.S. G.P.O. Washington, D.C.
- Wood, C. A. 1980. Morphometric evolution of cinder cones. <u>Journal of Volcanology and Geothermal</u> Research 7: 387-413.
- Woodcock, A. H. 1974. Permafrost and climatology of a Hawaii volcano crater. <u>Artic and Alpine</u> Research **6**(1): 49-62.
- Woodcock, A. H. 1980. Hawaiian alpine lake level, rainfall trends, and spring flow. <u>Pacific Science</u> **34**(2): 195-209.
- Woodcock, A. H. 1987. Mountain breathing revisited the hyperventilation of a volcano cinder cone. American Meteorological Society Bulletin **68**(2): 125-130.
- Woodcock, A. H. and G. W. Groves 1969. Negative thermal gradient under alpine lake in Hawaii. <u>Deep</u>Sea Research **16**: 393-405.
- Woodcock, A. H. and I. Friedman 1980. Mountain breathing: preliminary studies of air-land interaction on Mauna Kea, Hawaii. <u>Shorter contributions to geophysics</u>, 1979.
- Woodcock, A. H., A. S. Furumoto, et al. 1970. Fossil ice in Hawaii? Nature 226(5248): 873.
- Woodcock, A. H., R. Meyer, et al. 1966. Deep layer of sediments in alpine lake in the tropical mid-Pacific. Science **154**: 647-648.
- Woodward, S. A., P. M. Vitousek, K. Matson, F. Hughes, K. Benvenuto and P. A. Matson. 1990. Use of the exotic tree Myrica faya by native and exotic birds in Hawaii Volcanoes National Park. <u>Pacific Science</u> 44(1): 88-93.
- Woodworth, B. L., C. T. Atkinson, D. A. LaPointe, P. J. Hart, C. S. Spiegel, E. J. Tweed, C. Henneman, J. LeBrun, T. Denette, R. DeMots, K. L. Kozar, D. Triglia, D. Lease, A. Gregor, T. Smith and D. Duffy. 2005. Host population persistence in the face of introduced vector-borne diseases: Hawaii amakihi and avian malaria. Proceedings of the National Academy of Sciences 102(5): 1531 1536.
- World Conservation Monitoring Centre. 1998. Dubautia arborea. Retrieved November 12, 2007, from www.iucnredlist.org.

- Wright, T. L. and D. A. Clague. 1989. Petrology of Hawaiian lava. E. L. Winterer, D. M. Hussong and R. W. Decker. Boulder, CO, Geological Society of America. **The Geology of North America, v. N:** p. 218-237; pls. 1A, 1B, 2A, 2B, 3A, and 5 (folded maps) in separate box include Hawaii.
- Wunderle, J. M., Jr. 1997. The role of animal seed dispersal in accelerating native forest regeneration on degraded tropical lands. <u>Forest Ecology and Management</u> 99: 223-235.
- Xu, G., F. A. Frey, et al. 2005. East Molokai and other Kea-trend volcanoes: magmatic processes and sources as they migrate away from the Hawaiian hot spot. <u>G</u>³ (<u>Geochemistry</u>, <u>Geophysics</u>, <u>Geosystems</u>) **v. 6**(no. Q05008): doi:10.1029/2004GC000830.
- Yang, H.-J., F. Frey, et al. 1999. Mineral chemistry of submarine lavas from Hilo Ridge, Hawaii: implications for magmatic processes within Hawaiian rift zones. Contributions to Mineralogy and Petrology v. 135: p. 355-372.
- Yorinks, N. and C. T. Atkinson. 2000. Effects of malaria on activity budgets of experimentally infected juvenile Apapane (Himatione sanguinea). <u>Auk</u> 117(3): 731-738.
- Zavaleta, E. S. 2002. It's often better to eradicate, but can we eradicate better? <u>Turning the tide: the eradication of invasive species</u>. C. R. Veitch and M. N. Clout. Gland, Switzerland and Cambridge, UK, IUCN SSC Invasive Species Specialist Group: 393-403.
- Ziegler, A. C. 2002. <u>Hawaiian Natural History, Ecology and Evolution</u>. Honolulu HI, University of Hawaii Press.
- Zimmerman, E. C. 1941. Argentine ants in Hawaii. <u>Proceedings of the Hawaiian Entomological Society</u> 11(1): 108.
- Zimmerman, E. C. 1958. *Insects of Hawaii. Vol 8. Lepidoptera: Pyraloidea*. Honolulu, HI, University of Hawaii Press.